

# Checklist and distribution of lady beetles (Coleoptera: Coccinellidae) in Iran

Abdolahi Mesbah, R.,<sup>a</sup>, Nozari, J.<sup>a\*</sup>, Allahyari, H.<sup>a</sup>, Zare Khormizi, M.<sup>b</sup>

<sup>a</sup>Department of Plant Protection, College of Agriculture and Natural Resources, University of Tehran, Karaj, Iran

<sup>b</sup>Department of Entomology, College of Agricultural Sciences, Shiraz Branch, Islamic Azad University, Shiraz, Iran

(Received: 22 October 2015; Accepted: 4 July 2016)

This paper is an attempt to summarize all previous literature of Coccinellidae records from Iran and their distribution in the world as well as in provinces of Iran. There are 8 subfamilies, 17 tribes, 41 genera and 142 species in Iran recorded up to now. The largest subfamily, tribe and genera in Iran were Scymninae, Scymnini and *Scymnus*, respectively. The most widely distributed species in Iran were *Hippodamia variegata* (Goeze), *Coccinella septempunctata* L., *Oenopia conglobata* L., *Chilocorus bipustulatus* L. and *Adalia bipunctata* L.

**Key words:** Checklist, distribution, Iran, lady beetles.

## INTRODUCTION

Family Coccinellidae (lady beetles or ladybirds) is the largest family in the newly recognized superfamily Coccoidea (Robertson et al. 2015), and in the previously joint Cucujooidea s.lat., with about 6000 species described worldwide (Iablokoff-Khanzorian, 1982; Vandenberg, 2002). They are well known as mostly beneficial insects, their larvae and adults feed on pests, especially on aphids, scale insects and mites (Dobzhansky, 1933). But some of them are mycophagous and phytophagous and can become pests. Some species are very sensitive to changes in the environment so they can offer a better overall picture about the health of ecosystems and they can be good indicator species (Iperti, 1999).

Iran is an ecologically diverse country which includes different climates and geographical zones in 31 provinces. Rich faunal assemblage is thus expected for the country. There are many articles, theses and books about faunistic studies and coccinellid records in Iran and in the world. This paper is an attempt to review all previous literature of coccinellid records from Iran and species distribution in the world and also in different provinces of Iran. So far, a checklist of Scymninae of Iran (Jafari et al., 2013) and a catalog of Palearctic Coccinellidae (Kovar, 2007), including Iranian Coccinellidae, have been published. Subfamilies Microweiseinae and Sticholotidinae in Iran were recently summarized (Biranvand et al. in press a). This paper is an up to date list of species including their distribution in the world and in different provinces of Iran.

## MATERIAL AND METHODS

A revised list of the Coccinellidae of Iran was generated by reviewing the relevant literature on Coccinellidae. According to the last classification of Coleoptera, Coccinellidae were classified in two subfamilies Microweiseinae and Coccinellinae (Bouchard et al., 2011) but we use the classification of Kovar (2007) modified by Nedvěd and Kovář (2012) in this paper. We implemented some comments and synonyms, which have been used wrongly as separate species in some Iranian

\*Corresponding Author: nozari@ut.ac.ir

©2016 FERDOWSI UNIVERSITY OF MASHHAD, IRAN

articles. This list incorporated the most recent systematic and nomenclature updates. The arrangement of the subfamilies, tribes and genera are alphabetical for convenience and do not reflect phylogenetic relationships. Geographical distribution in 31 provinces of Iran and in the world was extracted for each species. For the distributions in the world, classification of geographical regions in the book Catalogue of Palaearctic Coleoptera (Ivan Löbl and Ales Smetana, 2007) was used.

## RESULTS

By reviewing the relevant literature on Coccinellidae of Iran, it was recognized that there are 142 species classified in eight subfamilies, 17 tribes and 41 genera in Iran.

### Subfamily CHILOCORINAE

#### Tribe Chilocorini

##### Genus *Brumoides* E. A. Chapin, 1965

###### *B. adenensis* Fursch, 1987

Distribution in Iran: Recorded from Iran by Kovar (2007) without mention of province

Distribution in the world: Afrotropical: Ethiopia (Fursch, 1987); Palearctic: Iran , Saudi Arabia, United Arab Emirates, Yemen (Kovar, 2007)

##### Genus *Chilocorus* Leach, 1815

###### *C. bipustulatus* (Linnaeus, 1758)

Distribution in Iran: Lorestan (Mohammad poor et al., 2013), Yazd (Zare Khormizi et al., 2013), Kerman (Salehi et al., 2011), North Khorasan (Biranvand et al. in press b), Razavi Khorasan (Farahi & Sadeghi, 2009), Semnan (Dazyanian & Sahragard, 2000), Fars (Fallahzadeh et al., 2000), Tehran (Omid et al., 2000), Kohgiluyeh and Boyer-Ahmad (Saeedi, 1998), Hamadan (Sadeghi & Khanjani, 1998), Mazandaran (Maafi et al., 1998), Zanjan (Mohseni et al., 1998), Gilan (Hai-Zadeh et al., 1998), Alborz (Babmorad, 1993), Chaharmahal and Bakhtiari (Esfandiari et al., 2002), Isfahan (Jafari et al., 2002), Golestan (Monutazeri & Mossadegh, 1995), South Khorasan (Moadi & Mossadegh, 1995), Markazi (Ahmadi et al., 2012), Khuzestan (Mossadegh, 2012), West Azerbaijan (Parvizi et al., 1986), Ardebil (Lotfalizadeh, 2001), Kermanshah (Khosravi et al., 2014), East Azerbaijan & Kordestan (Borumand, 2000)

Distribution in the world: Afrotropical: Sudan (Yousof et al., 2013); Nearctic: USA (Gordon, 1985); wide distribution in Palearctic (Mader, 1955; Kovar, 2007) including Mongolia (Bielawski, 1975)

##### Genus *Exochomus* Redtenbacher, 1844

(= *Brumus* Mulsant, 1850, = *Anexochomus* Barovskij, 1922)

###### *E. bifasciatus* Barovskij, 1927

Distribution in Iran: Recorded from Iran by Kovar (2007) without mention of province.

Distribution in the world: Palearctic: China, Iran and Kazakhstan (Kovar, 2007).

###### *E. gebleri* (Weise, 1885)

Distribution in Iran: Lorestan (Jafari & Kamali, 2007), Golestan (Kovar, 1995).

Distribution in the world: Palearctic: Afghanistan, Iran and Turkey (Kovar, 2007).

###### *E. quadriguttatus* (A. Fleischer, 1900)

Comment: Synonyme with *E. illaesicollis* (Roubal, 1927)

Distribution in Iran: Kerman (Yazdani & Ebrahimi, 1993)

Distribution in the world: wide distribution in Palearctic (Kovar, 2007)

***E. octosignatus* (Gebler 1830)**

Distribution in Iran: Chaharmahal and Bakhtiari (Bagheri & Mossadegh, 1995), South Khorasan (Mudi & Mosadegh, 2000), Gilan (Hai-Zadeh et al., 1998), Kerman (Yazdani & Ebrahimi, 1993), Razavi Khorasan (Yaghmaee & Kharaji, 1995), Isfahan & Alborz (Vodjdani, 1965), Lorestan (Ghanbari Kohyani et al., 2013), Hamadan (Hamedi & Moharrampour, 2012), West Azerbaijan & Qom & Tehran & Sistan and Baluchestan (Borumand, 2000)

Distribution in the world: Palearctic: Afghanistan, Albania, Armenia, France, Iran, Iraq, Italy, Russia, Kazakhstan, Kyrgyzstan, Mongolia, Tadzhikistan, Turkey, Turkmenistan, and Uzbekistan (Kovar, 2007)

***E. quadripustulatus* (Linnaeus, 1758)**

Distribution in Iran: Lorestan (Jafari & Kamali, 2007), Kerman (Salehi et al., 2011), Fars (Lotfalizade et al., 2000), Qazvin (Mohammadbeigi, 2000), Kohgiluyeh and Boyer-Ahmad (Saeedi, 1998), Zanjan (Mohseni et al., 1998), Isfahan (Bagheri & Emami, 1998), Chaharmahal and Bakhtiari (Esfandiari et al., 2002), Khuzestan (Mosadegh & Aleosfoor, 2004), Razavi Khorasan (Yaghmaee & Kharaji, 1995), North Khorasan (Biranvand et al. in press b), Mazandaran (Vodjdani, 1965), Yazd (Zare Khormizi et al., 2013) West Azerbaijan (Parvizi et al., 1986), Tehran (Abdi et al., 2012), East Azerbaijan & Qom & Ilam (Borumand, 2000)

Distribution in the world: Nearctic: USA (Gordon, 1985); Oriental: India (Poorani, 2004); Palearctic: wide distribution in western Palearctic (Mader, 1955; Kovar, 2007) and Russia (Poorani, 2004)

***E. undulatus* (Weise, 1878)**

Distribution in Iran: Lorestan (Mohammad poor et al., 2013), Kerman (Salehi et al., 2011), Fars (Fallahzadeh et al., 2000), Kohgiluyeh and Boyer-Ahmad (Saeedi, 1998), Isfahan (Bagheri & Emami, 1998), Chaharmahal and Bakhtiari (Esfandiari et al., 2002), Tehran (Ghanbari et al., 2012), Kermanshah (Gholami moghadam et al., 2014), Razavi Khorasan (Farahi & Sadeghi, 2009), South Khorasan (Moadi & Mossadegh, 1995), Markazi (Ahmadi et al., 2012)

Distribution in the world: Palearctic: Afghanistan, Albania, Egypt, Georgia, Iran, Iraq, Israel, Lebanon, Syria, (Kovar, 2007), Uzbekistan (Mader, 1955)

***E. kiritschenkoi* Barovskij, 1927**

Distribution in Iran: Recorded from Iran by Mader (1955) without mention of province.

Distribution in the world: Afghanistan, Iran, Turkmenistan (Mader, 1955); Kazakhstan, Kyrgyzstan, Mongolia and Tadzhikistan (Kovar, 2007)

**Genus *Parexochomus* Barovskij, 1922*****P. pubescens* (Kuster, 1848)**

Distribution in Iran: Lorestan (Ansari pour & Shakarami, 2011), Mazandaran (Montazeri & Mossadegh, 1995), South Khorasan (Moadi & Mossadegh, 1995), Kerman (Jafari et al., 2011), Khuzestan (Afshari, et al., 2000), Fars (Al-e-Mansoor & Ahmadi, 1993), Isfahan (Haghshenas et al., 2004), Sistan and Baluchestan (Modarres Najaf Abadi et al., 2008), Tehran (Samin & Shojai, 2013)

Distribution in the world: Afrotropical: Africa (Raimundo & Van Harten, 2000); Oriental: India (Raimundo et al., 2008); Palearctic: Afganistan, Algeria, Egypt, France, Germany, Iran, Iraq, Israeal, Italy, Lebanon, Morocco, Pakistan, Saudi Arabia, Spain, Tunisia (Kovar, 2007), Transcaucasia (Mader, 1955) and Turkey (Efil et al., 2010)

***P. melanocephalus* (Zoubkoff, 1833)**

Distribution in Iran: Lorestan (Jafari & Kamali, 2007), Kerman (Salehi et al., 2011), Razavi Khorasan (Yaghmaee & Kharaji, 1995), Chaharmahal and Bakhtiari (Bagheri & Mossadegh, 1995)

Distribution in the world: Palearctic: Albania, Armenia, Bulgaria, Georgia, Iran, Kazakhstan, Kyrgyzstan, Tadzhikistan, Turkey, Turkmenistan, Uzbekistan (Kovar, 2007), South Russia and Transcaucasia (Mader, 1955)

### ***P. nigripennis* (Erichson, 1843)**

Distribution in Iran: Yazd (Zare Khormizi et al., 2013), Kerman (Salehi et al., 2011), Semnan (Dazyanian & Sahragard, 2000), Razavi Khorasan (Kalantari & Sadeghi, 2000), Gilan (Hai-Zadeh et al., 1998), Fars (Al-e-Mansoor & Ahmadi, 1993), Golestan (Monutazeri & Mossadegh, 1995), South Khorasan (Moadi & Mossadegh, 1995), Khuzestan (Mosadegh & Aleosfoor, 2004), Tehran (Samin & Shojaei, 2013), Isfahan & Ilam & Khuzestan & Sistan and Baluchestan (Borumand, 2000)

Distribution in the world: Afrotropical: Angola, Chad, Tangier (Morocco), Luxor (Egypt), Tripoli (Lybia), Nubien (Sudan) (Fursch, 1961); Oriental: Northwestern India (Poorani, 2004); Palearctic: Afghanistan, Algeria, Canary Islands, Egypt, Italy, Iraq, Israel, Lebanon, Morocco, Pakistan, Portugal, Saudi Arabia, Spain, Syria, Tunisia, United Arab Emirates, (Kovar, 2007), Iran (Fursch, 1961; Kovar, 2007), Armenia, Azerbaijan, Georgia, Malta, Turkey (De Jong, 2011) and Caspian surrounding (Mader, 1955)

### ***P. nigromaculatus* Goeze, 1777**

Comment: This species is very similar to *P. flavipes* (Thunb, 1784) and because of this *P. flavipes* has been reported wrongly in some Iranian papers

Distribution in Iran: Lorestan (Jafari & Kamali, 2007), Kerman (Jafari et al., 2011), Razavi Khorasan (Farahi & Sadeghi, 2009), Gilan (Nazari et al., 2000), Hamadan (Sadeghi & Khanjani, 1998), Zanjan (Mohseni et al., 1998), Isfahan (Bagheri & Emami, 1998), Fars (Ahmadi & Sarafrazi, 1993), West Azerbaijan (Khalil Aria & Malkeshi, 2002), Chaharmahal and Bakhtiari (Esfandiari et al., 2002), Khuzestan (Mosadegh & Aleosfoor, 2004), Golestan (Monutazeri & Mossadegh, 1995), South Khorasan (Moadi & Mossadegh, 1995), North Khorasan (Biranvand et al. in press b), Markazi (Ahmadi et al., 2012), Tehran (Ghanbari et al., 2012), Kermanshah (Khosravi et al., 2014), East Azerbaijan & Ilam & Mazandaran (Borumand, 2000), Alborz (Vodjdani, 1965), Ardebil (Lotfalizadeh, 2001)

Distribution in the world: Nearctic: USA (Gordon, 1985); Palearctic: wide distribution in Palearctic (Mader, 1955; Kovar, 2007)

## **Subfamily COCCIDULINAE**

### **Tribe Coccidulini**

#### **Genus *Coccidula* Kugelann, 1798**

#### ***C. lithophiloides* Reitter, 1890**

Distribution in Iran: Kermanshah & Fars & Lorestan (Fursch, 1977)

Distribution in the world: Palearctic: Armenia (Kovar, 2007), Iran (Fursch, 1977; Kovar, 2007)

#### ***C. rufa* Herbst, 1783**

Distribution in Iran: Recorded from Iran by Kovar (2007) without mention of province

Distribution in the world: wide distribution in western Palearctic, Iran (Kovar, 2007)

#### **Genus *Lindorus* Casey, 1899**

#### ***L. lophantae* Blaisdell, 1892**

Comment: Also placed in *Rhyzobius* J.F. Stephens, 1832

Distribution in Iran: Introduced. Recorded from Iran by Kovar (2007) without mention of province

Distribution in the world: Australian: Australia, New Zealand (Canepari, 2011); Nearctic: introduced to USA (Raimundo & Van Harten, 2000); Neotropical: introduced to Chile (Canepari, 2011); Afrotopical: South and West Africa (Raimundo & Van Harten, 2000), Palearctic: Albania, Algeria, Canary Islands, Croatia, France, Germany, Italy, Iran, Morocco, Spain, Tunisia, Turkey, Ukraine, Yugoslavia (Serbia, Montenegro) (Kovar, 2007)

#### **Genus *Cryptolaemus* Mulsant, 1853**

##### ***C. montrouzieri* Mulsant, 1853**

Distribution in Iran: Fars & Gilan & Golestan & Lorestan & Mazandaran (Jafari et al., 2013), Kerman (Jafari et al., 2011), Khuzestan (Mossadegh et al., 2008), Markazi (Ahmadi et al., 2012)

Distribution in the world: Afrotropical: introduced to South Africa (Mader, 1955); Australian: Australia (Canepari, 2011), Tasmania, introduced to Hawaii and New Zealand (Mader, 1955); Nearctic: introduced to USA (Gordon, 1985); Neotropical: introduced to Chile, French Guyana (Canepari, 2011); Oriental: introduced to India (Poorani, 2004), Java (Mader, 1955); Palearctic: introduced to China, Egypt, France, Germany, Iran, Italy, Spain, (Kovar, 2007)

#### **Tribe Tetrabrachini**

##### **Genus *Tetrabrachys* Kapur, 1948**

##### ***T. bipustulatus* Barovskij, 1909**

Distribution in Iran: Lorestan (Iablokoff-Khnzorian, 1974)

Distribution in the world: Palearctic: Afganistan, Iran, Turkmenistan (Kovar, 2007)

##### ***T. coloratus* Fursch, 1960**

Distribution in Iran: Golestan (Fursch, 1960)

Distribution in the world: Palearctic: Iran (Fursch, 1960)

'

##### ***T. major* Crotch, 1874**

Distribution in Iran: Kordestan (Mader, 1955)

Distribution in the world: Palearctic: Iran (Mader, 1955; Kovar, 2007)

##### ***T. minimus* Barovskij, 1910**

Distribution in Iran: Lorestan (Mader, 1955; Iablokoff-Khnzorian, 1974)

Distribution in the world: Palearctic: Iran (Mader, 1955; Kovar, 2007)

##### ***T. naviauxi* Duverger, 1983**

Distribution in Iran: Lorestan, Mazandaran, Isfahan (Duverger, 1983)

Distribution in the world: Palearctic: Iran (Duverger, 1983)

##### ***T. pallidus* Crotch, 1874**

Distribution in Iran: Recorded from Iran by Mader (1955) without mention of province

Distribution in the world: Palearctic: Iran (Mader, 1955; Kovar, 2007)

##### ***T. pellucidus* Barovskij, 1909**

Distribution in Iran: Lorestan (Iablokoff-Khnzorian, 1974)

Distribution in the world: Palearctic: Iran (Mader, 1955; Kovar, 2007)

##### ***T. smarti* Iablokoff-Khnzorian, 1974**

Distribution in Iran: Recorded from Iran by Kovar (2007) without mention of province

Distribution in the world: Palearctic: Iran (Kovar, 2007)

***T. zarudnyi* Barovskij, 1909**

Distribution in Iran: Khorasan Razavi (Tablokoff-Khnzorian, 1974)

Distribution in the world: Palearctic: Iran (Tablokoff-Khnzorian, 1974; Mader, 1955)

**Subfamily COCCINELLINAE****Tribe Halyziini****Genus *Halyzia* Mulsant, 1846*****H. sedecimguttata* (Linnaeus, 1758)**

Distribution in Iran: Gilan & Mazandaran (Borumand, 2000)

Distribution in the world: wide distribution in Palearctic (Kovar, 2007), including Japan, Mongolia (Beilawski, 1968)

**Genus *Psyllobora* Dejean, 1836*****P. bisoctonotata* (Mulstantz, 1850)**

Distribution in Iran: Hormozgan & Fars (Borumand, 2000)

Distribution in the world: Palearctic: Afganistan, United Arab Emirates, Egypt, Iraq, Iran, Israel, Jordan, Pakistan, Saudi Arabia, Syria, Yemen (Kovar, 2007); Afrotropical (Kovar, 2007); Oriental: India (Kovar, 2007)

***P. vigintiduopunctata* (Linnaeus, 1758)**

Distribution in Iran: Hamadan (Sadeghi & Khanjani, 1998), Zanjan (Mohseni et al., 1998), Chaharmahal and Bakhtiari (Esfandiari et al., 2002), West Azerbaijan (Alizadeh & Malkeshi, 2010), North Khorasan (Biranvand et al. in press b), South Khorasan (Moadi & Mossadegh, 1995), Alborz (Vodjdani, 1965), Lorestan (Mohammad poor et al., 2013), Kerman (Koohpayezadeh & Mossadegh, 1991), Mazandaran (Montazeri & Mossadegh, 1995), Razavi Khorasan (Yaghmae & Kharazi Pakdel, 1995), Gilan (Haji Zade et al., 1998), Fars (Fallahzadeh et al., 2000), East Azerbaijan (Sadaghian et al., 2000), Isfahan (Nematollahi & Bagheri, 2002), Ardebil (Lotfalizadeh, 2001), Tehran (Borumand, 2000)

Distribution in the world: wide distribution in Palearctic (Kovar, 2007)

**Genus *Vibidia* Mulsant, 1846*****V. duodecimguttata* (Poda in Neuhaus, 1761)**

Distribution in Iran: East Azerbaijan (Mohammadi et al., 2013), West Azerbaijan (Parvizi et al., 1986), Gilan (Borumand, 2000), North Khorasan (Biranvand et al. in press b)

Distribution in the world: wide distribution in Palearctic (Kovar, 2007), including Mongolia (Beilawski, 1975)

**Tribe Coccinellini****Genus *Adalia* Mulsant, 1846*****A. bipunctata* (Linnaeus, 1758)**

Comment: Synonyme with *A. fasciatopunctata* (Faldermann, 1835)

Distribution in Iran: Yazd (Zare Khormizi et al., 2013), Kerman (Jafari et al., 2011), Lorestan (Mohammad poor et al., 2013), Chaharmahal and Bakhtiari (Bagheri & Mossedegh, 1995), Ardebil (Razmjo & Hajizadeh, 2000), West Azerbaijan (Akbarzadeh & Rezvani, 2000), Semnan (Dazyanian & Sahragard, 2000), North Khorasan (Biranvand et al. in press b), Razavi Khorasan (Kalantari & Sadeghi, 2000) Qazvin (Mohammadbeigi, 2000), Fars (Fallahzadeh et al., 2000), Hamadan (Ahmadi, 2000), Kohgiluyeh and Boyer-Ahmad (Saeedi, 1998), South Khorasan (Malkeshi, et al., 1998), Gilan (Hai-Zadeh et al., 1998), Golestan (Sadeghi et al., 2002), Isfahan (Jafari et al., 2002), Khuzestan

(Mosadegh & Aleosfoor, 2004), Sistan and Baluchestan (Modarres Najaf Abadi et al., 2008), Mazandaran & Tehran & Hormozgan (Vodjdani, 1965), Kermanshah (Gholami moghadam et al., 2014), Zanjan (Mohseni et al., 1998), East Azerbaijan & Qom & Kordestan (Borumand, 2000) Distribution in the world: Afrotropical: Middle of Africa (Bielawski, 1975); Nearctic: USA (Gordon, 1985) and Canada (Larson, 2013); Neotropical: introduced to Chile (Rebolledo et al., 2007); Oriental: India, (Poorani, 2004); Palearctic: wide distribution in Palearctic (Kovar, 2007), including Mongolia (Bielawski, 1975) and China (Poorani, 2004)

#### ***A. decempunctata* (Linnaeus, 1758)**

Distribution in Iran: Kerman (Jafari at al., 2011), Lorestan (Mohammad poor et al., 2013), Chaharmahal and Bakhtiari (Bagheri & Mossedegh, 1995), Golestan (Monutazeri & Mossadegh, 1995), North Khorasan (Biranvand et al. in press b), South Khorasan (Moadi & Mossadegh, 1995), Alborz (Vodjdani, 1965), Markazi (Ahmadi et al., 2012), Isfahan (Fatemi, 1982), Hamadan (Akhavan et al., 2013), Ilam & Tehran & Gilan (Borumand, 2000)

Distribution in the world: wide distribution in western Palearctic (Kovar, 2007), including Mongolia (Bielawski, 1975); Oriental: India (Lal & Kanakavalli, 1960)

#### ***A. tetraspilota* (Hope, 1831)**

Distribution in Iran: Lorestan (Biranvand et al., 2014), Razavi Khorasan (Borumand, 2000)

Distribution in the world: Oriental: India (Poorani, 2004) and Nepal (Poorani, 2004; Ashfaque, 2012); Palearctic: Afghanistan, China, Iran, Tadzhikistan, Turkmenistan and Uzbekistan, (Kovar, 2007)

#### **Genus *Aphidecta* Weise, 1893**

##### ***A. oblitterata* (Linnaeus, 1758)**

Distribution in Iran: Lorestan (Jafari et al., 2008)

Distribution in the world: Nearctic: Canada, USA (Gordon, 1985); Palearctic: wide distribution in western Palearctic (Kovar, 2007)

#### **Genus *Calvia* Mulsant, 1850**

##### ***C. punctata* (Mulsant, 1853)**

Distribution in Iran: Recorded from Iran by Kovar (2007) without mention of province

Distribution in the world: Oriental: India, Nepal (Ashfaque, 2012); Palearctic: Afganistan, Iran, Tadzhikistan and Uzbekistan (Kovar, 2007) and Paskistan (Poorani, 2004)

#### ***C. quatuordecimguttata* (Linnaeus, 1758)**

Distribution in Iran: Fars (Aliche & Yazdani, 1989), Kerman (Koohpayezadeh & Mossadegh, 1991)

Distribution in the world: Nearctic: USA (Bielawski, 1975) and Canada (Larson, 2013); Oriental: India, Bhutan and Nepal (Poorani, 2004); Palearctic: wide distribution in Palearctic (Kovar, 2007) including Mongolia (Bielawski, 1975)

#### **Genus *Ceratomegilla* Crotch, 1873**

##### ***C. undecimnotata* D. H. Schneider, 1792**

Distribution in Iran: Recorded from Iran by Kovar (2007) without mention of province

Distribution in the world: Palearctic: wide distribution in western Palearctic (Kovar, 2007), Caucasus, Western Siberia, Asia Minor (Iablokoff-Khnzorian, 1982)

#### **Genus *Coccinella* Linnaeus, 1758**

##### ***C. iranica* Dobzhansky, 1926**

Distribution in Iran: Mazandaran & Tehran (Naim, 1971)

Distribution in the world: Palearctic: Iran, Turkey (Kovar, 2007) and Pakistan (Ashfaque *et al.*, 2013)

#### ***C. magnopunctata* Rybakow, 1889**

Distribution in Iran: Razavi Khorasan (Farahi & Sadeghi, 2009)

Distribution in the world: Oriental: Mongolia, Nepal, Tibet (Poorani, 2004); Palearctic: China, Mongolia (Kovar, 2007), Iran (Farahi & Sadeghi, 2009); Oriental: India

#### ***C. septempunctata* Linnaeus, 1758**

Distribution in Iran: All parts of Iran

Distribution in the world: Nearctic: USA (Gordon, 1985), Canada (Larson, 2013); Oreintal: India (Bielawski, 1968; Canepari, 2011), Sri Lanka (Ashfaque, 2012), Pakistan (Poorani, 2004); Palearctic: wide distribution in Palearctic; Afrotropical (Kovar, 2007)

#### ***C. undecimpunctata* Linnaeus, 1758**

Distribution in Iran: Lorestan (Ansari pour & Shakarmi, 2011), Mazandaran (Fursch, 1977; Maafi *et al.*, 1998), Yazd (Zare Khormizi *et al.*, 2013), Ardebil (Razmjoo & Hajizadeh, 2000), Semnan (Dazyanian & Sahragard, 2000), Khuzestan (Bagheri & Mosadegh, 2000), East Azerbaijan (Sadaghian *et al.*, 2000), Hamadan (Sadeghi & Khanjani, 1998), Gilan (Hai-Zadeh *et al.*, 1998), Fars (Al-e-Mansoor & Ahmadi, 1993), Kohgiluyeh and Boyer-Ahmad (Saeedi, 2002), Sistan and Baluchestan (Modarres Najaf Abadi *et al.*, 2008), Golestan (Monutazeri & Mossadegh, 1995), Alborz & Tehran (Vodjdani, 1965), Isfahan (Vodjdani, 1965), Kerman (Salehi *et al.*, 2011), North Khorasan (Biranvand *et al.* in press b), South Khorasan (Kalantari & Sadeghi, 2000), West Azerbaijan (Parvizi *et al.*, 1986), Ilam (Borumand, 2000)

Distribution in the world: Australian: Australia (Poorani, 2004), New Zealand (Raimundo & Van Harten, 2000); Nearctic: Canada, USA (Gordon, 1985); Oriental: India, Nepal (Poorani, 2004); Palearctic: wide distribution in western Palearctic (Kovar, 2007), Mongolia (Bielawski, 1975) and Pakistan (Poorani, 2004); North Africa (Poorani, 2004)

#### **Genus *Harmonia* Mulsant, 1846**

##### ***H. quadripunctata* (Pontoppidan, 1763)**

Distribution in Iran: North Khorasan (Hamidi, 2013)

Distribution in the world: wide distribution in western Palearctic (Kovar, 2007)

#### **Genus *Hippodamia* Chevrolat, 1836**

##### ***H. septemmaculata* (DeGeer, 1775)**

Comment: Synonyme with *H. heidemanni* (Meier, 1897)

Distribution in Iran: Tehran (Borumand, 2000)

Distribution in the world: wide distribution in western Palearctic, Iran (Kovar, 2007)

##### ***H. variegata* (Goeze, 1777)**

Comment: Synonyme with *H. novemnotata* Haworth, 1912; also placed in *Adonia* Mulsant, 1846

Distribution in Iran: All parts of Iran

Distribution in the world: Afrotropical: Kenya (Canepari, 2007), Central Africa (Raimundo & Van Harten, 2000), Eastern Africa (Ashfaque, 2012); Neotropical: introduced to Chile (Rebolledo *et al.*, 2007); Oriental: India (Bielawski, 1968; Canepari, 2007), Nepal (Ashfaque, 2012); Palearctic: wide distribution in western Palearctic, Iran (Kovar, 2007), Tibet, China (Ashfaque, 2012) and Pakistan (Poorani, 2004)

***H. tredecimpunctata* (Linnaeus, 1758)**

Distribution in Iran: Kohgiluyeh and Boyer-Ahmad (Saeedi, 1998), Chaharmahal and Bakhtiari (Bagheri & Mossadegh, 1995), Isfahan (Fatemi, 1982), West Azerbaijan (Parvizi et al., 1986), Fars & Gilan & Mazandaran (Borumand, 2000)

Distribution in the world: Nearctic: Canada (Belicek, 1976); Oriental: Nepal, Tibet, India (Poorani, 2004); Palearctic: wide distribution in western Palearctic, Iran (Kovar, 2007)

**Genus *Menochilus* Timberlake, 1943*****M. sexmaculatus* (Fabricius, 1781)**

Comment: Also placed in *Cheilomenes* Mulsant, 1850

Distribution in Iran: East Azerbaijan (Sadaghian et al. 2000), Kerman (Koohpayezadeh & Mossadegh, 1991), Razavi Khorasan (Kalantari & Sadeghi, 2000), Fars (Mosadegh & Aleosfoor, 2004), Lorestan (Ghanbari Kohyani et al., 2013), Hormozgan & Sistan and Baluchestan & Tehran (Borumand, 2000)

Distribution in the world: Australian: Australia (Poorani, 2004); Oriental: Bhutan, China, Ceylon, Formosa, Malaysia, Micronesia, New Guinea, Philippines (Sasaji, 1971), India (Lal & Kanakavalli, 1960) Indonesia (Korschefsky, 1932), Timor (Alves & Raimundo, 1971), Bangladesh, Myanmar, Vietnam (Ashfaque, 2012), Nepal, (Kovar, 2007), Sri Lanka (Poorani, 2004); Palearctic: Afghanistan, Iran, Japan, Oman, Pakistan, United Arab Emirates; Afrotropical (Kovar, 2007)

**Genus *Oenopia* Mulsant, 1850*****O. conglobata* (Linnaeus, 1758)**

Comment: including two subspecies *O. conglobata contaminata* Ménétries, 1849 & *Oenopia conglobata conglobata* (Linnaeus, 1758)

Distribution in Iran: Lorestan (Farahbakhsh, 1961), Kerman (Salehi et al., 2011), Fars (Yazdani, 1990), Gorgan (Montazeri & Mossadegh, 1995), Kermanshah (Jalilvand et al., 2014) Mazandaran (Maafi et al., 1998), Razavi Khorasan (Farahi & Sadeghi, 2009), Semnan (Dazyanian & Sahragard, 2000), Qazvin (Mohammadbeigi, 2000), Alborz (Babmorad et al., 2000), Kohgiluyeh and Boyer-Ahmad (Saeedi, 1998), Hamadan (Sadeghi & Khanjani, 1998), North Khorasan (Biranvand et al. in press b), South Khorasan (Malkeshi, et al., 1998), Gilan (Hai-Zadeh et al., 1998), Tehran (Omid et al., 1993), Chaharmahal and Bakhtiari (Esfandiari et al., 2002), Isfahan (Jafari et al., 2002), Khuzestan (Mosadegh & Aleosfoor, 2004), Golestan (Monutazeri & Mossadegh, 1995), Zanjan (Mohseni et al., 1998), East Azerbaijan & Kordestan & Hormozgan (Vodjdani, 1965), West Azerbaijan (Parvizi et al., 1986), Ardebil (Lotfalizadeh, 2001), Qom (Borumand, 2000)

Distribution in the world: Nearctic: North America (Ashfaque, 2012); Oriental: India (Sicard, 1907); Palearctic: wide distribution in Palearctic (Kovar, 2007), Northern China (Ashfaque, 2012) and Pakistan (Poorani, 2004)

***O. montana* Savoyskaya, 1969**

Distribution in Iran: Recorded from Iran by Kovar (2007) without mention of province

Distribution in the world: Palearctic: Iran & Kazakhstan (Kovar, 2007)

***O. oncina* (Olivier, 1808)**

Distribution in Iran: Lorestan (Ansari pour & Shakarami, 2011), Yazd (Zare Khormizi et al., 2013), Kerman (Jafari et al., 2011), Razavi Khorasan (Kalantari & Sadeghi, 2000), Isfahan (Seifollahi et al., 2000), Fars (Al-e-Mansoor & Ahmadi, 1993), West Azerbaijan (Khalil Aria & Malkeshi, 2002), Chaharmahal and Bakhtiari (Esfandiari et al., 2002), Khuzestan (Mosadegh & Aleosfoor, 2004), South Khorasan (Moadi & Mossadegh, 1995), Markazi (Ahmadi et al., 2012), Kermanshah (Gholami

Moghadam et al., 2014), Hamadan (Akhavan et al., 2013), Qom & Ilam & Mazandaran & Sistan and Baluchestan & Tehran (Borumand, 2000)

Distribution in the world: wide distribution in western Palearctic, Iran (Kovar, 2007)

### **Genus *Propylea* Mulsant, 1846**

#### ***P. quatuordecimpunctata* (Linnaeus, 1758)**

Distribution in Iran: Alborz (Farahbakhsh, 1961), Lorestan (Mohammad poor et al., 2013), Chaharmahal and Bakhtiari (Bagheri & Mossedegh, 1995), North Khorasan (Biranvand et al. in press b), Razavi Khorasan (Farahi & Sadeghi, 2009), Kerman (Jafari et al., 2011), Ardebil (Razmjoh & Hajizadeh, 2000), Kohgiluyeh and Boyer-Ahmad (Saeedi, 1998), Hamadan (Sadeghi & Khanjani, 1998), Mazandaran (Maafi et al., 1998), Gilan (Hai-Zadeh et al., 1998), Golestan (Sadeghi et al., 2002), Isfahan (Haghshenas et al., 2004), Fars (Aliche & Yazdani, 1989), Khuzestan & Alborz & Hormozgan (Vodjdani, 1965), Tehran (Abdi et al., 2012), Qom & Semnan (Borumand, 2000)  
Distribution in the world: Nearctic: USA & Canada (Hoebelke & Wheeler, 1996); Oriental: India (Ashfaque, 2012), Bangladesh (Poorani, 2004); Palearctic: wide distribution in Palearctic (Kovar, 2007), including China (Ashfaque, 2012) and Pakistan (Poorani, 2004)

### **Tribe Tytthaspidini**

#### **Genus *Anisosticta* Dejean, 1837**

##### ***A. caucasica* A. Fleischer, 1900**

Distribution in Iran: Tehran (Fursch, 1977)

Distribution in the world: Palearctic: Armenia (Kovar, 2007) Iran (Fursch, 1977; Kovar, 2007)

##### ***A. bitriangularis* (Say, 1842)**

Distribution in Iran: Razavi Khorasan (Yaghmaee & Kharaji, 1995)

Distribution in the world: Nearctic: North America (Bielawski, 1958) and Canada (Larson, 2013); Palearctic: Iran (Kovar, 2007), Russia (Far East), Siberia (Bielawski, 1958)

##### ***A. novemdecimpunctata* (Linnaeus, 1758)**

Distribution in Iran: Alborz (Vodjdani, 1965)

Distribution in the world: wide distribution in western Palearctic, Iran (Kovar, 2007)

### **Genus *Bulaea* Mulsant, 1850**

#### ***B. lichatschovii* Hummel, 1827**

Distribution in Iran: South Khorasan (Moadi & Mossadegh, 1995), Alborz (Vodjdani, 1965), Isfahan (Fatemi, 1982), West Azerbaijan (Parvizi et al., 1986), Ardebil (Lotfalizadeh, 2001), East Azerbaijan & Bushehr & Fars & Hormozgan & Ilam & Kerman & Sistan and Baluchestan & Markazi (Borumand, 2000)

Distribution in the world: Afrotropical: Central Africa (Poorani, 2004); Oriental: India (Poorani, 2004); Palearctic: wide distribution in Palearctic (Kovar, 2007)

#### ***B. lividula* bocanderi Mulsant, 1850**

Distribution in Iran: Recorded from Iran by Kovar (2007) without mention of province

Distribution in the world: Palearctic: Algeria, Egypt, Morocco, United Arab Emirates, Iraq, Israel, Jordan, Pakistan, Saudi Arabia, Syria, Yemen, Iran; Afrotropical (Kovar, 2007)

### **Genus *Coccinula* Dobzhansky, 1925**

#### ***C. elegantula* (Weise, 1890)**

Distribution in Iran: Yazd (Zare Khormizi et al., 2013), Isfahan (Bagheri & Emami, 1998), Razavi Khorasan (Yaghmaee & Kharaji, 1995), Chaharmahal and Bakhtiari (Bagheri & Mossadegh, 1995), Lorestan (Ansari pour & Shakarami, 2011), Kermanshah (Jalilvand et al., 2014)

Distribution in the world: Palearctic: Russia, Iran, Kazakhstan, Mongolia, Uzbekistan, Kyrgyzstan, Tadzhikistan (Kovar, 2007), Mongolia and Turkestan (Duverger, 1983)

#### ***C. redimita* (Weise 1885)**

Distribution in Iran: Chaharmahal and Bakhtiari (Bagheri & Mossedegh, 1995), Lorestan (Ghanbari Kohyani et al., 2013), Tehran (Borumand, 2000)

Distribution in the world: Palearctic: Afghanistan, Iran, Tadzhikistan, Uzbekistan (Kovar, 2007), Turkey & Pakistan (Ashfaque, 2012) and Mongolia (Bielawski, 1975)

#### ***C. sinuatomarginata* Faldermann, 1837**

Distribution in Iran: Tehran (Borumand, 2000)

Distribution in the world: wide distribution in Palearctic (Kovar, 2007)

#### **Genus *Tytthaspis* Crotch, 1874**

##### ***T. sedecimpunctata* (Linnaeus, 1758)**

Distribution in Iran: Lorestan (Mohammad Poor et al., 2013)

Distribution in the world: wide distribution in Palearctic (Kovar, 2007) including Mongolia (Bielawski, 1975)

#### **Subfamily EPILACHINAE**

##### **Tribe Epilachini Mulsant, 1846**

##### **Genus *Chnootriba* Chevrolat, 1837**

##### ***C. elaterii* P. Rossi, 1794**

Distribution in Iran: Bushehr & Isfahan & Fars & Gilan & Hormozgan & Ilam & Kermanshah & Khuzestan & Lorestan & Semnan & Tehran (Borumand, 2000), Golestan (Khormali, 2012), North Khorasan (Biranvand et al. in press b),

Distribution in the world: North Africa: Niger, Sudan, Senegal, Chad, Burkina Faso (Poorani, 2004); Oriental: India (Poorani, 2004); Palearctic: Algeria, Albania, Austria, Bosnia Herzegovina, Bulgaria, Croatia, Czech Republic, France, Great Britain, Germany, Hungary, Italy, Luxembourg, Morocco, Portugal, Romania, Spain, Slovakia, Switzerland, Turkey, Tunisia, Ukraine, (Kovar, 2007), Iran (Borumand, 2000), Turkey (Poorani, 2004)

##### **Genus *Cynegetis* Chevrolat, 1836**

##### ***C. syriaca* Mader, 1958**

Comment: Synonyme with *C. rapillyi* Duverger, 1983

Distribution in Iran: Kerman (Duverger, 1983)

Distribution in the world: Palearctic: Iran, Syria, Turkey (Kovar, 2007)

##### **Genus *Henosepilachna* Li, 1961**

##### ***H. argus* (Geoffroy, 1785)**

Distribution in Iran: Ilam (Borumand, 2000), Kermanshah (Modarrese aval, 1994)

Distribution in the world: palearctic: Armenia, Austria, Belgium, Bulgaria, Czech Republic, Great Britain, France, Germany, Hungary, Italy, Netherlands, Poland, Romania, Slovakia, Spain, Sweden, Yugoslavia, Algeria, Morocco, Lebanon, Tadzhikistan, Turkmenistan, Turkey, Uzbekistan (Kovar, 2007), Iran (Bagheri & Modarress Aval, 1994); Afrotropical (Kovar, 2007)

##### ***H. vigintiquatuorpunctata* (Linnaeus, 1758)**

Distribution in Iran: Recorded from Iran by Kovar (2007) without mention of province  
 Distribution in the world: wide distribution in Eastern Palearctic (Kovar, 2007)

### **Subfamily MICROWEISEINAE**

#### **Tribe Microweiseini Leng, 1920**

##### **Genus *Paracoelopterus* Normand, 1936 (=*Diloponis* Pope 1962)**

###### ***P. berytensis* J. Weise, 1884**

Comment: Synonyme with *P. fuerschi* Ahmadi & Yazdani 1993

Distribution in Iran: Fars (Ahmadi & Yazdani, 1993), Khorasan Razavi (Khandehroo, 2015)

Distribution in the world: Palearctic: Germany, Iran, Israel, Lebanon, Morocco, Tunisia, (Kovar, 2007), Syria (Mader, 1955)

### **Tribe Serangiini**

#### **Genus *Serangium* Blackburn, 1889**

##### ***S. montazerii* Fursch, 1995**

Distribution in Iran: Zanjan (Mohseni et al., 1998), Gilan (Hai-Zadeh et al., 1998), Golestan (Monutazeri & Mossadegh, 1995)

Distribution in the world: Palearctic: France, Georgia, Iran, Israel, Pakistan, Syria (Kovar, 2007), Oriental: India (?)

### **Subfamily Ortaliinae Mulsant, 1850**

#### **Tribe Noviini Mulsant, 1850**

##### **Genus *Rodolia* Mulsant, 1850**

###### ***R. cardinalis* Mulsant, 1850**

Distribution in Iran: Gilan (Hai-Zadeh et al., 1998), Khuzestan (Eslamizadeh & Barzkar, 2006), Fars (Alichi & Yazdani, 1989), Golestan (Monutazeri & Mossadegh, 1995), Mazandaran (Borumand, 2000)

Distribution in the world: Australian: Australia (Gordon, 1972); Afrotropical: introduced to North and South Africa (Gordon, 1972); Nearctic: introduced to USA (Gordon, 1985); Neotropical: introduced to South America (Gordon, 1972); Oriental: introduced to India (Lal & Kanakavalli, 1960); Palearctic: widely introduced in Palearctic (Kovar, 2007)

##### ***R. fausti* Weise, 1885**

Distribution in Iran: Razavi Khorasan (Yaghmaee & Kharaji, 1995)

Distribution in the world: Palearctic: Iran, Kazakhstan, Kyrgyzstan, Uzbekistan, (Kovar, 2007)

##### ***R. rufipennis* Pic, 1926**

Distribution in Iran: Recorded from Iran by Kovar (2007) without mention of province

Distribution in the world: Palearctic: Egypt, Iran and Libya (Kovar, 2007)

### **Subfamily SCYMNINAE**

#### **Tribe Diomini R. D. Gordon, 2000**

##### **Genus *Diomus* Mulsant, 1850**

###### ***D. rubidus* Motscholsky, 1837**

Distribution in Iran: Kerman (Koohpayezadeh & Mosadegh 1991), Kermanshah (Jalilvand et al., 2014)

Distribution in the world: wide distribution in western Palearctic, Iran (Kovar, 2007)

**Tribe Hyperaspidini****Genus *Hyperaspis* Chevrolat 1837*****H. concolor* (Suffrian, 1843)**

Distribution in Iran: Chaharmahal and Bakhtiari (Bagheri & Mosadegh 1995)

Distribution in the world: Palearctic: Austria, Belgium, Bulgaria, Czech Republic, Finland, France, Germany, Hungary, Italy, Poland, Romania, Slovakia, Sweden, Switzerland (Kovar, 2007), Iran (Bagheri & Mosadegh 1995)

***H. duvergeri* Fursch, 1985**

Distribution in Iran: Chaharmahal and Bakhtiari (Bagheri & Mosadegh 1995)

Distribution in the world: Palearctic: Algeria, Croatia, France, Hungary, Italy, Portugal, Spain, Morocco (Kovar, 2007), Iran (Bagheri & Mosadegh 1995)

***H. femorata* (Motschulsky, 1837)**

Distribution in Iran: Tehran (Borumand, 2000)

Distribution in the world: Palearctic: Armenia, Azerbaijan, Bulgaria, Russia, Tadzhikistan, Turkmenistan, Ukraine, (Kovar, 2007), Iran (Borumand, 2000)

***H. histeroides* Faldermann, 1837**

Distribution in Iran: Recorded from Iran by Kovar (2007) without mention of province

Distribution in the world: Palearctic: Armenia, Azerbaijan, Iran, Kazakhstan (Kovar, 2007)

***H. marmottani* Fairmaire, 1868**

Distribution in Iran: Razavi Khorasan (Yaghmaee & Kharazipakdel 1995)

Distribution in the world: Algeria, Tunisia (Sicard, 1907; Kovar, 2007), Egypt, Iran, Libya, (Kovar, 2007)

***H. persica* Duverger, 1983**

Distribution in Iran: Kerman (Duverger, 1983), Khorasan (Moadi & Mossadegh, 1995)

Distribution in the world: Palearctic: Iran (Duverger, 1983)

***H. polita* Weise, 1885**

Distribution in Iran: Golestan & Kerman & South Khorasan & Razavi Khorasan & Khuzestan (Jafari et al., 2013), Fars (Fahhahzade et al., 2006), Kermanshah (Jalilvand et al., 2014: Khosravi et al., 2014)

Distribution in the world: Afrotropical: Chad, Ethiopia, Sudan, Somalia, Kenya (Duverger, 1983); Oriental: India (Iablokoff-Khnzorian, 1971); Palearctic: Afghanistan, Cyprus, Egypt, Iran, Israel, Syria, Tadzhikistan, Turkmenistan, Turkey, (Kovar, 2007), Coastal areas of the eastern Mediterranean region, Lebanon, Pakistan, (Alizadeh et al., 2013)

***H. pumila* (Sharafme 1987)**

Distribution in Iran: Tehran (Naeem 1971), South Khorasan (Moadi & Mossadegh, 1995)

Distribution in the world: Afrotropical: Cameron, Congo, Erythrea, Guinea, Sudan, (Duverger, 1983), Senegal (Sicard A., 1907; Iablokoff-Khnzorian, 1971; Duverger, 1983); Palearctic: Algeria, Egypt, Libya, Iran (Duverger, 1983), Yemen, Arabia (Raimundo et al., 2006), Egypt (Mader, 1955), Lybia, Tunisia (Iablokoff-Khnzorian, 1971)

***H. quadrimaculata* (Redtenbacher, 1843)**

Distribution in Iran: Lorestan, Mazandaran, Isfahan (Duverger, 1983), (Ansaripour 2012), Alborz (Ansari pour, 2012)

Distribution in the world: Palearctic: Albania, Austria, Belgium, Bulgaria, Czech Republic, Germany, Hungary, Romania, Slovakia, Lebanon, Turkey, Syria (Kovar, 2007), Iran (Duverger, 1983)

### ***H. reppensis* (Herbst, 1783)**

Distribution in Iran: Yazd (Zare et al., 2014)

Distribution in the world: wide distribution in western Palearctic (Kovar, 2007) and Iran (Zare et al., 2014)

### ***H. syriaca* Weise, 1885**

Distribution in Iran: Mazandaran (Duverger, 1983), Chaharmahal and Bakhtiari (Bagheri & Mosadegh 1995)

Distribution in the world: Palearctic: Israel, Syria, Lebanon, Turkey, Iran (Kovar, 2007), Lebanon (Fursch, 1977)

### ***H. transversoguttata* J. Weise, 1878**

Distribution in Iran: Kerman, Khuzestan (Duverger, 1983)

Distribution in the world: Palearctic: Armenia, Iran, Kazakhstan, Uzbekistan, Mongolia, Tadzhikistan, Turkmenistan, Kyrgyzstan, China (Kovar, 2007), Afghanistan, Azerbaijan, (Iablokoff-Khnzorian, 1971)

### ***H. vinciguerrae* Capra, 1929**

Distribution in Iran: Kuzestan (Alizadeh et al., 2013)

Distribution in the world: Afrotropical: Northern and eastern Africa (Raimundo & Van Harten, 2000); Palearctic: Libya, Saudi Arabia (Raimundo & Van Harten, 2000) Yemen (Iablokoff-Khnzorian, 1971; Raimundo & Van Harten, 2000), Iran (Alizadeh et al., 2013)

### **Tribe Platynaspidini**

#### **Genus *Platynaspis* Redtenbacher, 1844**

##### ***P. luteorubra* (Goeze, 1777)**

Distribution in Iran: Lorestan (Jafari & Kamali, 2007), South Khorasan (Moadi & Mossadegh, 1995), Ardebil (Lotfalizadeh, 2001), Fars & Tehran (Borumand, 2000)

Distribution in the world: wide distribution in western Palearctic, Iran (Kovar, 2007)

### **Tribe Scymnini**

#### **Genus *Clitostethus* Weise, 1885**

##### ***C. arcuatus* (Rossi, 1794)**

Distribution in Iran: Chaharmahal and Bakhtiari & Fars & Golestan & Kerman & South Khorasan & Gilan (Jafari et al., 2013) Ardebil (Lotfalizadeh, 2001), North Khorasan (Biranvand et al. in press b)

Distribution in the world: wide distribution in western Palearctic, Iran (Kovar, 2007); Neotropical: introduced to Chile (Canepari, 2011); Afrotropical (Kovar, 2007)

#### **Genus *Nephus* Mulsant, 1846**

##### ***N. arcuatus* Kapur**

Distribution in Iran: Khuzestan (Mossadegh et al., 2012)

Distribution in the world: Afrotropical: Togo (Raimundo et al., 2008); Palearctic: Iran, Syria, Yemen (Kovar, 2007), Saudi Arabia (Raimundo & Van Harten, 2000), United Arab Emirates (Raimundo et al., 2008)

***N. biguttatus* Mulsant, 1850**

Distribution in Iran: Golestan & South Khorasan (Jafari et al., 2013) Chaharmahal and Bakhtiari (Esfandiari et al., 2002), Kerman (Kouhpayezadeh & Mossadegh, 1991)

Distribution in the world: Turkey (Kaydan et al., 2006)

***N. bipunctatus* (Kugelann, 1794)**

Distribution in Iran: Fars (Ahmadi & Yazdani 1993), Kerman (Kouhpayezadeh Esfahani & Mosadegh 1991), Semnan (Dazyanian & Sahragard, 2000), Zanjan (Mohseni et al., 1998), Gilan (Hajizadeh et al., 1998), Isfahan (Fatemi, 1982)

Distribution in the world: wide distribution in western Palearctic (Kovar, 2007), Iran (Kouhpayezadeh Esfahani & Mosadegh 1991), Japan (Mader, 1955)

***N. bisignatus* Fursch, 1984**

Distribution in Iran: Razavi Khorasan (Yaghmaee & Kharazipakdel 1995)

Distribution in the world: Palearctic: Austria, Czech Republic, Germany, Hungary, Sweden (Kovar, 2007), Iran (Yaghmaee & Kharazipakdel 1995)

***N. caneparii* Fursch & Uygun, 1980**

Distribution in Iran: Recorded from Iran by Kovar (2007) without mention of province

Distribution in the world: Palearctic: Iran and Turkey (Kovar, 2007)

***N. caucasicus* Weise, 1929**

Distribution in Iran: Tehran (Jafari et al., 2013)

Distribution in the world: Palearctic: Armenia, Iran (Kovar, 2007), Kaukasus (Mader, 1955)

***N. conjunctus* Wollaston, 1870**

Comment: Synonyme with *N. includens* Kirsch, 1871

Distribution in Iran: Khuzestan (Asadeh & Mosaddegh, 1991; Mossadegh & Kocheili, 1993), Fars (Fallahzadeh et al., 2004)

Distribution in the world: Palearctic: Canary Islands, Egypt, Greece, Israel, Saudi Arabia, Portugal, Turkey, Yemen (Raimundo et al., 2006.), Iran (Asadeh & Mosaddegh, 1991), Turkestan (Mader, 1955); Afrotropical: Cape Verde Islands, Gambia, Guinea Bissau, Ethiopia, (Raimundo et al., 2006)

***N. crucifer* A. Fleischer, 1900**

Distribution in Iran: Recorded from Iran by Kovar (2007) without mention of province

Distribution in the world: Palearctic: Iran, Kazakhstan, Kyrgyzstan, Saudi Arabia, Tadzhikistan, Turkmenistan, Uzbekistan (Kovar, 2007), Yemen (Raimundo & Van Harten, 2000); Afrotropical (Kovar, 2007)

***N. fenestratus* J. R. Sahlberg, 1913**

Distribution in Iran: Khuzestan (Asadeh & Mosaddegh, 1991, Mossadegh & Kocheili, 1993), Fars (Fallahzadeh & Hesami, 2004), Kermanshah (Jalilvand, 2011)

Distribution in the world: Palearctic: United Arab Emirates, Cyprus, Egypt, Israel, Saudi Arabia, Iran (Kovar, 2007), Syria (Raimundo & Van Harten, 2000), United Arab Emirates (Raimundo et al., 2008); Afrotropical (Kovar, 2007)

***N. hiekei* Fursch, 1987**

Distribution in Iran: Fars (Falahzade, et al. 2003)

Distribution in the world: Palearctic: United Arab Emirates, Algeria, Egypt, Greece, Iran, Iraq, Italy, Lebanon, Morocco, Saudi Arabia, Spain, Tunisia, Turkey (Kovar, 2007), Syria, Yemen (Raimundo et al., 2008), Israel (Fursch, 1967), Portugal (Raimundo, A. 1992)

***N. jucundus* Fursch, 1977**

Distribution in Iran: Azerbaijan (Fursch, 1977)

Distribution in the world: Palearctic: Iran (Fursch, 1977)

***N. levaillanti* Mulsant, 1850**

Comment: Synonyme with *S. levaillantis* Mulsant, 1850

Distribution in Iran: Fars & Kerman & Razavi Khorasan & Khuzestan (Jafari et al., 2013)

Distribution in the world: Afrotropical: Rwanda (Fursch, 1997); Oriental: India (Canepari, 1986); Palearctic: Afghanistan, China, Egypt, Germany, Iran, Israel, Italy, Japan, Jordan, Russia, Lebanon, Pakistan (Kovar, 2007)

***N. nigricans* (Weise, 1879)**

Distribution in Iran: Lorestan (Biranvand, 2014)

Distribution in the world: Albania, Crotia, France, Greece, Italy, Turkey (Kovar, 2007), Iran (Biranvand, 2014)

***N. quadrimaculatus* (Herbst, 1783)**

Distribution in Iran: Kerman (Salehi et al., 2011), Kermanshah (Gholami Moghadam et al., 2014), Gilan & Golestan & Mazandaran (Jafari et al., 2013)

Distribution in the world: wide distribution in Palearctic (Kovar, 2007)

***N. ulrichi* Fursch, 1977**

Distribution in Iran: Chaharmahal and Bakhtiari (Bagheri & Mosadegh, 1995), Khuzestan (Mosadegh & Aleosfoor, 2004), Fars (Fahhhazade et al., 2006), Kermanshah (Jalilvand et al., 2014).

Distribution in the world: Palearctic: Greece, Italy, Portugal, Spain (Kovar, 2007), Iran (Bagheri & Mosadegh, 1995)

**Genus *Scymniscus* Dobrzhanskiy, 1925??*****S. biflammulatus* (Motschulsky, 1837)-**

Distribution in Iran: Khuzestan (Bagheri & Mosadegh, 1995) Lorestan (Biranvand, 2014), Kermanshah (Jalilvand et al., 2014: Khosravi et al., 2014)

Distribution in the world: Palearctic: Afghanistan, Armenia, Georgia, Iran, Turkmenistan, Uzbekistan (Kovar, 2007)

***S. uguttatus* Mulsant, 1850**

Comment: Synonyme with *S. quadriguttatus* Feischer, 1900

Distribution in Iran: Recorded from Iran by Kovar (2007) without mention of province

Distribution in the world: Palearctic: Afghanistan, Iran, Kazakhstan (Kovar, 2007)

***S. elbursi* Fursch, 1981**

Distribution in Iran: Alborz (Karaj) (Fursch, 1981), Kerman (Salehi et al., 2011)

Distribution in the world: Palearctic: Iran (Fursch, 1981)

**Genus *Scymnus* Kugelann, 1794**

***S. apetzi* Mulsant, 1846**

Distribution in Iran: West Azerbaijan & Chaharmahal and Bakhtiari & Fars & Gilan & Golestan & Razavi Khorasan & Lorestan & Tehran (Jafari et al., 2013), Mazandaran (Maafi et al., 1998), South Khorasan (Malkeshi, et al., 1998), Isfahan (Bagheri & Emami, 1998), Khuzestan (Mossadegh, 2012), Hamadan (Akhavan et al., 2013), North Khorasan (Biranvand et al. in press b)

Distribution in the world: wide distribution in western Palearctic, Iran (Kovar, 2007)

***S. arabicus* Fursch, 1989**

Distribution in Iran: Fars & Gilan & Lorestan (Jafari et al., 2013)

Distribution in the world: Palearctic: Saudi Arabia (Kovar, 2007), Iran (Jafari et al., 2013)

***S. argutus* Mulsant, 1850**

Distribution in Iran: Kerman (Jafari et al., 2013), Chaharmahal and Bakhtiari (Bagheri & Mossadegh, 1995)

Distribution in the world: Palearctic: Afghanistan, Albania, Armenia, Georgia, Iran, Iraq, Israel, Turkmenistan, Uzbekistan (Kovar, 2007)

***S. auritus* (Thunberg, 1795)**

Distribution in Iran: Chaharmahal and Bakhtiari & Fars & Yazd & Gilan (Jafari et al., 2013), Kerman (Salehi et al., 2011)

Distribution in the world: wide distribution in Palearctic (Kovar, 2007)

***S. ebneri* J. Weise, 1926**

Distribution in Iran: Recorded from Iran by Kovar (2007) without mention of province  
Distribution in the world: Palearctic: United Arab Emirates, Egypt, Iran; Afrotropical (Kovar, 2007)

***S. cedricolus* Fursch, 1977**

Distribution in Iran: Recorded from Iran by Kovar (2007) without mention of province

Distribution in the world: Palearctic: Iran, Lebanon, Turkey (Kovar, 2007)

***S. flagellisiphonatus* (Fursch, 1970)**

Distribution in Iran: North Khorasan (Hamidi, 2013)

Distribution in the world: wide distribution in Palearctic (Kovar, 2007)

***S. flavigollis* Redtenbacher, 1843**

Distribution in Iran: Chaharmahal and Bakhtiari & Fars & Gilan & Isfahan & Razavi Khorasan & Khuzestan & Lorestan (Jafari et al., 2013), Kerman (Salehi et al., 2011), Semnan (Dazyanian & Sahragard, 2000), Kermanshah (Bagheri & Emami, 1998), Tehran & East Azerbaijan (Fursch, 1977), Markazi (Ahmadi et al., 2012)

Distribution in the world: Palearctic: Algeria, France, Italy, Morocco, Tunisia (Kovar, 2007), Iran (Fursch, 1977), Syria (Mader, 1955), Lebanon (Fursch, 1977)

***S. fraxini* Mulsant, 1850**

Distribution in Iran: Recorded from Iran by Kovar (2007) without mention of province

Distribution in the world: wide distribution in western Palearctic (Kovar, 2007)

***S. frontalis* Fabricius, 1787**

Comment: Synonyme with *S. bimaculatus* (Thunberg, 1795)

Distribution in Iran: West Azerbaijan & Isfahan & Razavi Khorasan & Lorestan & Tehran & Gilan & Golestan (Jafari et al., 2013), Kermanshah (Gholami moghadam et al., 2014)

Distribution in the world: wide distribution in Palearctic (Kovar, 2007)

***S. impexus* Mulsant, 1850**

Distribution in Iran: West Azerbaijan (Parvizi et al., 1987), Razavi Khorasan (Kalantari & Sadeghi, 2000)

Distribution in the world: Nearctic: Canada, USA (Gordon, 1985); Oriental: India (Poorani, 2004); Palearctic: Austria, China, Czech Republic, France, Germany, Italy, Poland, Slovakia, Spain (Kovar, 2007), Iran (Parvizi et al., 1987), North Africa (Mader, 1955)

***S. indrihensis* Weise, 1929**

Distribution in Iran: South Khorasan (Moadi & Mossadegh, 1995)

Distribution in the world: wide distribution in Palearctic (Kovar, 2007), Iran (Moadi & Mossadegh, 1995), Turkestan (Mader, 1955), Mongolia (Bielawski, 1968)

***S. latemaculatus* Motschulsky, 1858**

Distribution in Iran: unknown

Distribution in the world: Palearctic: Afghanistan, China, Iran, Pakistan, (Kovar, 2007); Oriental: India, Sri Lanka, Taiwan, Thailand (Poorani, 2004)

***S. limbatus* Stephens, 1832a**

Distribution in Iran: Recorded from Iran by Kovar (2007) without mention of province

Distribution in the world: wide distribution in western Palearctic (Kovar, 2007)

***S. manipulus* Fursch & Kreissl, 1967**

Distribution in Iran: Recorded from Iran by Kovar (2007) without mention of province

Distribution in the world: Palearctic: China, Iran, Mongolia, Turkmenistan (Kovar, 2007)

***S. marginicollis* Mannerheim, 1843**

Comment: this species has been wrongly reported from Iran by Behdad (1988)

***S. marinus* (Mulsant, 1850)**

Comment: Synonyme with *S. mediterraneus* Iablokoff-Khnzorian, 1972 and *S. minimus* (O. F. Müller, 1776)

Distribution in Iran: Lorestan (Jafari et al., 2013)

Distribution in the world: Palearctic: Algeria, Canary Islands, Egypt, France, Italy, Morocco, Maderia Archipelago, Portugal, Spain, Tunisia (Kovar, 2007), Iran (Jafari et al., 2013), Syria (Capra, 1976)

***S. mongolicus* Weise, 1890**

Distribution in Iran: Kerman & South Khorasan & Khuzestan (Jafari et al., 2013), Isfahan (Bagheri & Emami, 1998), Fars (Mosadegh & Aleosfoor, 2004)

Distribution in the world: Palearctic: China, Mongolia (Kovar, 2007), Iran (Jafari et al., 2013)

***S. nigrinus* Kugellan, 1794**

Distribution in Iran: Isfahan (Jafari et al., 2013), Ardebil (Lotfalizadeh, 2001)

Distribution in the world: wide distribution in western Palearctic (Kovar, 2007), Iran (Jafari et al., 2013)

***S. nubilus* (Mulsant, 1850)**

Distribution in Iran: Hormozgan (Fallahzadeh et. al., 2013), Lorestan (Biranvand, 2014)

Distribution in the world: Afrotropical: Rwanda (Fursch, 1997); Oriental: India, Myanmar, Sri Lanka, Bangladesh (Poorani, 2004), Nepal (Ashfaque, 2012); Palearctic: wide distribution in western Palearctic (Kovar, 2007)

***S. pallipes* Mulsant, 1850**

Distribution in Iran: Chaharmahal and Bakhtiari & Golestan & Razavi Khorasan & Lorestan (Jafari et al., 2013), Kerman (Jafari at al., 2011), Isfahan (Bagheri & Emami, 1998), North Khorasan (Biranvand et al. in press b)

Distribution in the world: Palearctic: China, Iran, Kazakhstan, Russia, Tadzhikistan, Turkmenistan, Uzbekistan (Kovar, 2007), Iran (Jafari at al., 2011)

***S. pharaonis* Motschulsky, 1851**

Comment: Synonyme with *S. araraticus* (ablokoff-Khnzorian, 1969)

Distribution in Iran: Chaharmahal and Bakhtiari & Kerman (Jafari et al., 2013), Gilan (Hai-Zadeh et al., 1998), Lorestan (Jafari et al., 2008), Fars (Yazdani & Ahmadi, 1991), Kermanshah (Fursch, 1977)

Distribution in the world: Palearctic: Armenia, Egypt, France, Germany, Iraq, Jordan, Turkey (Kovar, 2007), Iran (Fursch, 1977; Kovar, 2007), Lebanon (Fursch, 1977)

***S. magnomaculatus* Fursch, 1958**

Comment: Synonyme with *S. quadriguttatus* (Capra, 19274)

Distribution in Iran: Fars & Gilan & Golestan & Razavi Khorasan (Jafari et al., 2013), Sistan and Baluchestan (Modarres Najaf Abadi et al., 2008), Ardebil (Lotfalizadeh, 2001)

Distribution in the world: Palearctic: Albania, Armenia, Austria, Azerbaijan, Bosnia Herzegovina, Bulgaria, Croatia, France, Greece, Italy, Macedonia, Romania, Slovenia, Turkey (Kovar, 2007), Iran (Jafari et al., 2013)

***S. rubromaculatus* Goeze, 1777**

Distribution in Iran: Fars & Gilan (Jafari et al., 2013), Semnan (Dazyanian & Sahragard, 2000), Isfahan (Seifollahi et al., 2000), Ardebil (Lotfalizadeh, 2001), Kermanshah (Gholami moghadam et al., 2014), Lorestan (Biranvand, 2014)

Distribution in the world: wide distribution in western Palearctic (Kovar, 2007); Afrotropical (Kovar, 2007)

***S. subvillosus* Goeze, 1777**

Distribution in Iran: West Azerbaijan & Fars & Isfahan & Kerman & Razavi Khorasan & Lorestan & Markazi (Jafari et al., 2013), Mazandaran (Maafi et al., 1998), Ardebil (Lotfalizadeh, 2001)

Distribution in the world: wide distribution in western Palearctic, Iran (Kovar, 2007); Afrotropical (Kovar, 2007)

***S. suffrianioides* Sahlberg, 1913**

Including two subsp.: *S. suffrianioides apetzioides* Capra & Fursch, 1967 and *S. suffrianioides suffrianioides* J. R. Sahlberg, 1913

Comment: *S. suffrianioides* Sahlberg, 1913 is synonym with *S. pallipediformis* Günther, 1958

Distribution in Iran: Isfahan & Razavi Khorasan & Lorestan & Tehran (Jafari et al., 2013) Markazi (Ahmadi et al., 2012), Kermanshah, Yazd (Zare et al., 2014)

Distribution in the world: Palearctic: Austria, France, Germany, Iraq, Israel, Italy, Jordan, Lebanon, Poland, Romania, Russia, Slovakia, Syria, Turkey (Kovar, 2007), Iran (Borumand, 2000)

### ***S. syriacus* (Marseul, 1898)**

Distribution in Iran: Chaharmahal and Bakhtiari & Fars & Gilan & Golestan & Kerman & South Khorasan & Razavi Khorasan & Khuzestan & Lorestan (Jafari et al., 2013), Yazd (Zare Khormizi et al., 2013), Semnan (Dazyanian & Sahragard, 2000), Zanjan (Mohseni et al., 1998), Isfahan (Bagheri & Emami, 1998), Ardebil (Lotfalizadeh, 2002), Markazi (Ahmadi et al., 2012), Hamadan (Akhavan et al., 2013), Tehran (Samin & Shojaei, 2013), North Khorasan (Biranvand et al. in press b)

Distribution in the world: Egypt, Cyprus, Iran, Iraq, Israel, Jordan, Lebanon, Syria (Kovar, 2007), Libya, Saudi Arabia, Turkey, Yemen (Alizadeh et al., 2013)

### ***S. testaceus* Motschulsky, 1837**

Distribution in Iran: Lorestan (Biranvand, 2014)

Distribution in the world: Palearctic: Albania, Kyrgyzstan, Tadzhikistan (Kovar, 2007), Iran (Biranvand, 2014)

### **Tribe Stethorini**

#### **Genus *Stethorus* Weise, 1885**

##### ***S. gilvifrons* Mulsant, 1850**

Distribution in Iran: Alborz (Farahbakhsh, 1961), Chaharmahal and Bakhtiari & Fars & Gilan & Golestan & South Khorasan & Razavi Khorasan & Khuzestan & Lorestan & Tehran & Isfahan (Jafari et al., 2013), Ardebil (Razmjoo & Hajizadeh, 2000), Hamadan (Eghbalian et al., 2008), Kerman (Koohpayezadeh & Mossadegh, 1991), Yazd (Zare et al., 2014), North Khorasan (Biranvand et al. in press b)

Distribution in the world: Oriental: India (Mader, 1955; Lal & Kanakavalli, 1960); Palearctic: Afghanistan, Armenia, Azerbaijan, Bulgaria, Croatia, Cyprus, Egypt, France, Greece, Iran, Iraq, Italy, Lebanon, (Kovar, 2007), Pakistan (Ashfaque, 2012), Turkey (Efil et al., 2010), Saudi Arabia (Raimundo & Van Harten, 2000)

### ***S. pusillus* (Herbst, 1797)**

Comment: Synonyme with *S. punctillum* (Weise, 1891)

Distribution in Iran: West Azerbaijan & Chaharmahal and Bakhtiari & Gilan & Isfahan & Kerman & South Khorasan & Razavi Khorasan & Khuzestan & Lorestan & Tehran (Jafari et al., 2013), Fars (Aliche & Yazdani, 1989), Alborz (Vodjani, 1965), Lorestan (Biranvand, 2014)

Distribution in the world: wide distribution in Palearctic (Kovar, 2007); Nearctic: USA (Gordon, 1985), Canada (Larson, 2013)

### ***S. siphonulus* (Kapur, 1948)**

Distribution in Iran: Khuzestan & Lorestan (Jafari et al., 2013)

Distribution in the world: Palearctic: China (Kovar, 2007), Iran (Jafari et al., 2013)

### **Subfamily STICHOLOTIDINAE**

#### **Tribe Sticholotidini**

##### **Genus *Coelopterus* Mulsant & Rey, 1852**

##### ***C. salinus* Mulsant & Rey, 1852**

Distribution in Iran: Recorded from Iran by Kovar (2007) without mention of province

Distribution in the world: Afrotropical: Somalia (Plaza, 1986); Palearctic: Algeria France, Iran, Italy, Morocco, Portugal, Spain, Tunisia, (Kovar, 2007), Syria (Mader, 1955)

**Genus *Pharoscymnus* Bedel, 1906**

***P. angohranensis* Duverger, 1983**

Distribution in Iran: Angothran (Duverger, 1983)

Distribution in the world: Iran (Duverger, 1983)

***P. arabicus* Fursch, 1979**

Distribution in Iran: Recorded from Iran by Kovar (2007) without mention of province

Distribution in the world: United Arab Emirates, Iran, Saudi Arabia (Kovar, 2007)

***P. brunneosignatus* Marder, 1949**

Distribution in Iran: North Khorasan (Biranvand et al. in press b), Razavi Khorasan (Ebrahimi et al. 2012)

Distribution in the world: Palearctic: China (Bielawski, 1975; Kovar, 2007), Mongolia (Kovar, 2007) and Iran (Ebrahimi et al., 2012, 2014)

***P. fleischeri* J. (Weise, 1883)**

Distribution in Iran: Recorded from Iran by Kovar (2007) without mention of province

Distribution in the world: Palearctic: Germany, Turkey (Kovar, 2007)

***P. flexibilis* (Mulstant, 1853)**

Distribution in Iran: Fars (Borumand, 2000)

Distribution in the world: Oriental: India, Ceylon (Korschefsky, 1931); Palearctic: Afghanistan, Pakistan, Iran (Kovar, 2007), Yemen, Oman (Raimundo & Van Harten, 2000)

***P. ovoideus* Sicard, 1907**

Distribution in Iran: Lorestan (Jafari & Kamali, 2007), Gilan (Hai-Zadeh et al., 1998), Fars (Lachinani & Ahmadi, 1993), Tehran (Abdi et al., 2012)

Distribution in the world: Palearctic: Algeria, Morocco, Tunisia, Jordan, Syria, Iran (Kovar, 2007), United Arab Emirates (Raimundo et al., 2008); Afrotropical (Kovar, 2007)

***P. pharoides* (Marsuel, 1868)**

Distribution in Iran: Fars (Ahmadi et al., 1993), Chaharmahal and Bakhtiari (Bagheri & Mossadegh, 1995)

Distribution in the world: Palearctic: Egypt, Israel, Syria, Libya, Saudi Arabia, Turkey, Iran; Afrotropical (Kovar, 2007)

***P. setulosus* (Chevrolat, 1861)**

Distribution in Iran: Kerman (Salehi et al., 2011), Fars (Yazdani & Ahmadi, 1992)

Distribution in the world: Palearctic: Algeria, United Arab Emirates, Egypt, Iran, Israel, Jordan Lebanon, Morocco, Saudi Arabia, Spain, Tunisia (Kovar, 2007).

**DISCUSSION**

According to the species distribution, the most widespread in Iran were the following species:

*Hippodamia variegata* and *Coccinella septempunctata* (recorded from all provinces of Iran), *Adalia bipunctata* (recorded from 26 provinces), *Oenopia conglobata* (recorded from 25 provinces), *Chilocorus bipustulatus* (recorded from 24 provinces).

Most species had limited distribution (recorded from less than 10 provinces).

The number of species recorded for each province is as follows:

Fars: (49), Lorestan: (47), Kerman: (44), Razavi Khorasan: (40), Tehran: (37), Isfahan: (36), Gilan: (34), Chaharmahal and Bakhtiari: (32), Khuzestan: (29), South Khorasan: (26), Golestan: (26), Kermanshah: (24), Mazandaran: (23), North Khorasan: (19), West Azerbaijan: (19), Ardebil: (18), Alborz: (17), Hamadan: (15), Yazd: (15), East Azerbaijan: (13), Semnan: (13), Markazi : (13), Ilam: (12), Sistan and Baluchestan: (11), Zanjan: (11), Kohgiluyeh and Boyer-Ahmad: (10), Hormozgan: (10), Qom: (9), Kordestan: (5), Qazvin: (5), Bushehr: (2)

The most species richness was belonging to provinces Fars, Lorestan and Kerman. Many scientists have recorded many ladybirds from the mentioned provinces and also they have rich and variable environments. On the other hand the least species richness was belonging to provinces Kordestan, Qazvin and Bushehr. Few scientists have recorded ladybirds from these provinces. They are small provinces and also Bushehr has a quite uniform environment.

According to the result of this study, the most species rich subfamily in Iran was Scymninae by almost 75 recorded species. The largest tribe was Scymnini by almost 60 recorded species and the largest genus was *Scymnus* by 28 recorded species.

#### Acknowledgments

We thank Dr Oldřich Nedvěd for comments that greatly improved the manuscript. We would also thank Dr Mostafa Mirzaei for providing some references.

#### LITERATURE CITED

- Abdi, A.R., Sadeghim, S.E., Talebi, A.A., Shojai, M., 2013. Coccinellid fauna of Chitgar Park and determination of dominant species. *Iranian Journal of Forest and Range Protection Research* 10(2), 146-164.
- Afshari, A., Mossadegh, M.S., Kamali, K., 2000. Spider mites, predators and their average feeding on *Tetranychus turkestanii* U. & N. in Khuzestan Province. Proceeding of the 14<sup>th</sup> Iranian Plant Protection Congress, 4-7 Sep 2000, Isfahan, Iran, pp 320.
- Ahmadi A., Yazdani, A., 1993. A new species of *Diloponis* from Iran. *Nachrichtenblatt Der Bayerischen Entomologen* 42, 30-32.
- Ahmadi, A., Jafari, R., Vafai, R., 2012. The faunistic survey of ladybird (Cole., Coccinellidae) in orchards and crops Arak and shrub. Proceeding of the 20<sup>th</sup> Iranian Plant Protection Congress, 25-28 Aug 2012, Shiraz, Iran, pp 139.
- Ahmadi, A.A., Aliche, M., Yazdani, A., 1993. A new record of *Pharoscymnus pharoides* Mars. From Iran. Proceeding of the 11<sup>th</sup> Iranian Plant Protection Congress, 27 Aug-1 Sep 1993, Rasht, Iran, pp 250.
- Ahmadi, A.A., Sarafrazi, A.M., 1993. Distribution in Iran and natural enemies of Russian Wheat aphid, *Diuraphis noxia* (Mordvilko) (Homoptera: Aphididae) in the Fars Province. Proceeding of the 11<sup>th</sup> Iranian Plant Protection Congress, 27 Aug-1 Sep 1993, Rasht, Iran, pp 1.
- Ahmadi, R., 2000. Natural enemies of Russian wheat aphid (*Diuraphis noxia* Mordvilko) in Hamedan province. Proceeding of the 14<sup>th</sup> Iranian Plant Protection Congress, Isfahan, 4-7 Sep 2000, Iran, pp 226.

Akandeh, M., Shishebor, P., Khajezadeh, Y., 2011. Sampling program and host preference of melon ladybeetle, *Epilachna chrysomelina* (F.) on some hosts of cucurbit family in Ahvaz, Southwest Iran. *Iranian Journal of Plant Protection Science* 42, 1-10.

Akbarzadeh Shoukat, G., Rezvani, A., 2000. Studies on apple aphids and their natural enemies in Urmia apple orchards. Proceeding of the 14<sup>th</sup> Iranian Plant Protection Congress, 4-7 Sep 2000, Isfahan, Iran, pp 259.

Akhavan, E., Jafari, R., Vafai, R., Afrogheh, S., 2013. Biodiversity and distribution in Iran of predaceous ladybird (Coleoptera: Coccinelliae). *International Research Journal of Applied and Basic Sciences* 5 (6): 705-709.

Al-e-Mansoor, H., Ahmadi, A.A., 1993. Natural enemies of cotton whitefly, *Bemisia tabaci* (Gennadius) in Fars province. Proceeding of the 11<sup>th</sup> Iranian Plant Protection Congress, 27 Aug-1 Sep 1993, Rasht, Iran, pp 106.

Ali, H.A., Abdul-Rassul, M.S., Swail, M.A., 1990. Systematic list of Coccinellidae recorded from Iraq. Bulletin of the Iraq Natural History Museum 8 (3), 45-51.

Aliche, M., Yazdani, A., 1989. Studies on the fauna of Coccinellidae in Fars. Proceeding of the 9<sup>th</sup> Iranian Plant Protection Congress, 8-13 Sep 1989, Mashhad, Iran, pp 9.

Alizadeh, E., Malkeshi, S.H., 2010. Study on the dinamism of dominant cereal aphid lady beetle species population in west Azerbaijan. Proceeding of the 19<sup>th</sup> Iranian Plant Protection Congress, 31 July-3 Aug 2010, Tehran, Iran, pp 57.

Alizadeh, M., Mossadegh, M., Esfandiari, M., 2013. Natural enemies of *Maconellicoccus hirsutus* (Green) (Hemiptera: Pseudococcidae) and their population fluctuations in Ahvaz, southwest of Iran. *Journal of Crop Protection* 2 (1), 13-21.

Alves, M.L.G, Raimundo, A.C., 1971. Coccinelídeos do Timor Portoguês. Garcia de Orta. *Lisboa* 19(1-4), 37-50.

Ansari Pour, A., 2012. Identification of some of the fauna ladybirds (Col.: Coccinellidae) in Alborz Province. Proceeding of the 20<sup>th</sup> Iranian Plant Protection Congress, 25-28 Aug 2012, Shiraz, Iran, pp 97.

Ansari Pour, A., Shakarami, J., 2011. Study of ladybirds (Col: Coccinellidae) in Khorramabad district and the first report of *Hyperaspis quadrimaculata* (Redtenbacher 1844) for Iranian fauna. *Life Science Journal* 8(3), 488-495.

Ansari Pour, A., Vafaei, S. R., Shahkarami, J., 2008. Study of fauna ladybirds (Col.: Coccinellidae) in Khorramabad district. *Journal of Entomological Research* 3 (2), 103-117.

Asadeh, G.A, Mosaddegh, M.S., 1991. The mealybugs, Pseudococcus spp. (Hemip.: Pseudococcidae) fauna of Khuzestan province. *The Scientific Journal of Agriculture* 16(1,2), 47-52.

Ashfaque, A., 2012. Taxonomic studies of family Coccinellidae (Coleoptera) of Gilgit-Baltistan, Pakistan. Department of plant protection. Faculty of crop protection sciences. The University of agriculture, Peshawar. 196p.

Ashfaque, M., Ullah, F., Rafi, M.A., 2013. Genus *Coccinella* (Coccinellidae: Coleoptera) from Gilgit-Baltistan with two new records from Pakistan. Sarhad Journal of Agriculture 29(2), 241-247.

Babmorad, M., 1993. Preliminary study of poplar insect fauna in Alborz Research Center in Karaj. Proceeding of the 11<sup>th</sup> Iranian Plant Protection Congress, 27 Aug-1 Sep 1993, Rasht, Iran, pp 238.

Babmorad, M., Bagheri, E., Yarmand, H., 2000. Identification of predacious insects of *Monosteria unicostata* in Karaj. Proceeding of the 14<sup>th</sup> Iranian Plant Protection Congress, 4-7 Sep 2000, Isfahan, Iran, pp 290.

Baghei, S., Mosadegh, M.S., 2000. Natural enemies of *Thrips tabaci* Lindeman. (Thysanoptera; Thripidae) of onion fields in Khozestan. Proceeding of the 14<sup>th</sup> Iranian Plant Protection Congress, 4-7 Sep 2000, Isfahan, Iran, pp 250.

Bagheri, M., Mosadegh, M.S., 1995. Fauna of Coccinellidae in Charmahal Bakhtiari province. Proceeding of the 12<sup>th</sup> Iranian Plant Protection Congress, 2-7 Sep 1995, Karaj, Iran, pp 308.

Bagheri, M.R., Emami, M.S., 1998. Introduction of ten new species of Coccinellidae for Isfahan Province. Proceeding of the 13<sup>th</sup> Iranian Plant Protection Congress, 3-8 Sep 1998, Karaj, Iran, pp 201.

Bagheri, M.R., Mossadegh, M.S., 1995. The faunistic studies of Coccinellidae in Chaharmahal Bakhtiari province. Proceeding of the 12<sup>th</sup> Iranian Plant Protection Congress, 2-7 Sep 1995, Karaj, Iran, pp 308.

Belicek, J., 1976. Coccinellidae of western Canada and Alaska with analyses of the transmontane zoogeographic relationships between the fauna of British Columbia and Alberta (Insecta Coleoptera Coccinellidae). *Quaestiones Entomologicae* 12, 283-409.

Bielawski, R., 1968. Ergebnisse der zoologischen Forschungen von Dr. Z. Kaszab in der Mongolei, 116. Coleoptera: Coccinellidae III. *Annales Zoologici* 26(4), 193-207.

Bielawski, R., 1975. Ergebnisse der zoologischen Forschungen von Dr. Z. Kaszab in der Mongolei Nr. 352. Coccinellidae V und VI. *Fragmenta Faunistica* 20(16), 247-271.

Biranvand, A., Jafari, R., Shakarmi, J., Vafaie-Shoushtari, R., Zare Khormizi, M., Ebarhimi, S., 2014. Faunistic study of ladybirds (Col.: Coccinellidae) in Chegeni region, Lorestan Province, Iran. Proceeding of the 21<sup>th</sup> Iranian Plant Protection Congress, 23-26 Aug 2014, Orumieh, Iran, pp 572.

Biranvand, A., Jafari, R., Zare, M., 2013. Polymorphism of *Hippodamia variegata* Goeze (Col.: Coccinellidae) in Lorestan Province. *Iran International Journal of Agriculture and Crop Sciences* 5 (21), 2655-2659.

Biranvand A., Nedvěd O., Tomaszewska W., Canepari C., Shakarami J., Fekrat L., Khormizi M.Z., 2016 a: An annotated checklist of Microweiseinae and Sticholotidini of Iran (Coleoptera, Coccinellidae). *ZooKeys*

Biranvand A, Shakarami J, Hamidi E, Fekrat L, Zare Khormizi M, Volf M, Nedvěd O, 2016 b: Ladybird beetles (Coleoptera: Coccinellidae) of North Khorasan province, their food plants, and new records for fauna of Iran. *Turkish Journal of Zoology*

Borumand, H., 2000. Insects of Iran: the list of Coleoptera in the insect collection of Plant Pests & Diseases Research Institute. Coleoptera (XXIV): Cucuoidea: Fam.(120) Coccinellidae. Plant Pests & Diseases Research institute, Insect taxonomy research department, Publ. No. 5, 44 pp.

Bouchard, P., Bousquet, Y., Davies, A.E., Alonso-Zarazaga, M.A., Lawrence, J.F., Lyal, C.H.C., Newton, A.F., Reid, C.A.M., Schmitt, M., Ślipiński, S.A., Smith, A.B.T., 2011. Family-group names in Coleoptera (Insecta). *ZooKeys* 88, 1–972.

Canepari, C., 1986. Sualcuni Coccinellidae dell'India e Nepal Settentrionale del Museo di Storia Naturale di Ginevra (Coleoptera: Coccinellidae). *Rivue Suisse de Zoologie* 93(1), 21–36

Canepari, C., 2007. I colleotteri Coccinellidi (Coleoptera: Coccinellidae) Conservazione Habitat Invertebrati 4, 213-219.

Canepari, C., 2011. Contribution too the Knowledge of Coccinellidae of Sardinia (Coleoptera), Conservazion Habitat Invertebratai 5, 501-516 .

Capra, F., 1976. Coccinellidae dell'Arcipelago Pontino (Coleoptera). 8 pp.

Chazeau, J., Fursch, H., Sasaji, H., 1989. Taxonomy of Coccinellidae. *Coccinella* 1, 6-8.

Chazeau, J., Fursch, H., Sasaji, H., 1990. Taxonomy of Coccinellidae (corrected version). *Coccinella* 2 (2), 4-6.

Chrif Smaili, M., Blenzar, A., Fursch, H., 2010. Entomologie faunistique – Faunistic Entomology 62(3), 103-107.

Davoodi Dehkordi, S., Sahragard, A., 2013. Functional Response of *Hippodamia variegata* (Coleoptera: Coccinellidae) to Different Densities of *Aphis gossypii* (Hemiptera: Aphididae) in an Open Patch Design. *Journal of Agricultural Science and Technology* 15(4), 651-659.

De Jong, Y.S.D.M., 2011. Fauna Europaea version 2.4. Web Service available online at <http://www.faunaeur.org>.

Dezianian, A., Sahragard, A., 2000. Investigation on natural enemies of the pistacia psyllid, *Agonoscena pistaciae* in Damgan-Iran. Proceeding of the 14<sup>th</sup> Iranian Plant Protection Congress, 4-7 Sep 2000, Isfahan, Iran, pp 270.

Dobzhansky, T., 1933. Geographical variation in lady-beetles. *American Naturalist* 67, 97-126.

Duverger, C., 1983. Contribution à la connaissance des Coccinellidae d'Iran. *Nouvelle Revue d'Entomologie XIII* (1), 73-93.

Ebrahimi, S., Karimi, J., Modarres-Awal, M., 2012. The first record of the ladybird, *Pharoscymnus brunneosignatus*, Mader, 1949 (Col., Coccinellidae) for Iran. Proceedings of the 20<sup>th</sup> Iranian Plant Protection Congress, 25-28 Aug 2012, Shiraz, Iran, pp 128.

Ebrahimi S, Modarres-Awal M, Fekrat L, Nedvěd O (2014) Two new records of ladybirds (Col.: Coccinellidae) for the Iranian beetle fauna. *Journal of Entomological Society of Iran* 34: 11–12.

Ebrahimzadeh, P., Mossadegh, M.S., 2004. The coccinellids and aphids fields in Khuzestan. Proceeding of the 16<sup>th</sup> Iranian Plant Protection Congress, 28 Aug-1 Sep 2004, Tabriz, Iran, pp 137.

Efil, L., Bayram, A., Ayaz, T., Senal, D., 2010. Coccinellidae species and their population changes in alfalfa field Akcakale country of Sanhurfa province and a new record *Exochomus pubescens* Küster from Turkey. *Bitki Koruma Bulteni* 50(3), 101-109.

Eghbalian, A.H., Khanjani, M., Pourmirza, A.A., 2008. Study on natural enemies of brown mite, *Bryobia scheuten* (Acari: Tetranychidae) in plum and cherry orchards in Hamedan province. Proceeding of the 18<sup>th</sup> Iranian Plant Protection Congress, 24-27 Aug 2008, Hamedan, Iran, pp 37.

Esfandiari, Radjab, G., Bagheri, M., Barari, H., 2002. Predator coccinellids on *Sphaerolecanium* Fonsc. (Hom. Coccidea) on almond and investigation on population changes of dominant species. Proceeding of the 15<sup>th</sup> Iranian Plant Protection Congress, 7-11 Sep 2002, Kermanshah, Iran, pp 91.

Eslamizadeh, R., Barzkar, M., 2006. Survey of stable of lady-bird *Rodalia (Vedalia) cardinalis* (Nab) in citrus orchard of Khuzestan Province. Proceeding of the 17<sup>th</sup> Iranian Plant Protection Congress, 2-5 Sep 2006, Tehran, Iran, pp 344.

Fallahzadeh, M., Shojai, M., Ostovan, H., Kamali, K., 2006. An investigation of the mealybug predators fauna in Fars Province-Iran. Proceeding of the 17<sup>th</sup> Iranian Plant Protection Congress, 2-5 Sep 2006, Tehran, Iran, pp 84.

Fallahzadeh, M., Abdimaleki, R., Saghaei, N., 2013. Contribution to the knowledge of the ladybird beetles (Coleoptera, Coccinellidae), redators of mealybugs (Hemiptera, Pseudococcidae) in Hormozgan province, Southern Iran. *Linzer Biologische Beitraege* 45(1), 673-679.

Fallahzadeh, M., Eghtedar, E., Ebrahimi, E., 2000. Natural enemies of olive psylla, *Euphyllura alivina* Costa in Fars province. Proceedings of the 14<sup>th</sup> Iranian Plant Protection Congress, 4-7 Sep 2000, Isfahan, Iran, pp 277.

Fallahzadeh, M., Hesami, S., 2004. Study of the natural enemies of *Maconellicoccus hirsutus* (Homoptera: Pseudococcidae) in Jahrom region of Fars province. Proceeding of the 16<sup>th</sup> Iranian Plant Protection Congress, 28 Aug-1 Sep 2004, Tabriz, Iran, 47 pp.

Fallahzadeh, M., Hesami, S., Fursch, H., 2003. Report of *Nephus (Sidis) hiekei* (Col.: Coccinellidae) in Iran. *Journal of Entomological Society of Iran* 22(2), 81-82.

Fallahzadeh, M., Hesami, SH., Alemansoor, H., 2004. Citrus whiteflies (Hemiptera: Aleyrodidae) and their natural enemies in Jahrom region of Fars province. Proceedings of the 16<sup>th</sup> Iranian Plant Protection Congress, 28 Aug-1 Sep 2004, Tabriz, Iran, pp 46.

Farahbakhsh, G., 1961. Checklist of important insects and other enemies of plants and agricultural products in Iran. Agriculture Plant Protection Organization 1-153.

Farahi, S., Sadeghi Namaghi, H., 2010. Report of *Coccinella magnopunctata* (Col.: Coccinellidae) from Iran. *Journal of Entomological Society of Iran* 30 (2), 79-80.

Farahi, S., Sadeghi Namaghi, H., 2009. Species diversity of aphids and lady beetles of wheat fields of Mashhad city (Razavi Khorasan Province). *Journal of Plant Protection* 23 (2), 89-95.

Fatemi, H., 1982. Fauna ladybirds of Isfahan province. *Journal of Plant pests and deseases* 50 (221), 21-25.

Fursch, H., 1960. Zwei neue Tetrabrachys-Arten (Col. Cocc.). Mitteilungen der Münchner Entomologischen Gesellschaft 50, 64-67.

Fursch, H., 1961. Revision der afrikanischen Arten um *Exochomus flavipes* Thunb. Col. Cocc. Entomologischen Arbeiten aus dem Museum G. Frey 12, 68-92.

Fursch, H., 1967. *Scymnus (Sidis) biekei* Fursch, 1987, als fiend von *Pseudococcus citri* Risso. *Entomoohaga* 12, 309-310.

Fursch, H., 1977. Coccinellidenausbeuten aus Libanon und dem Iran im Museum Genf mit Beschreibung neuer Scymnini-Arten. *Revue Suisse de Zoologie* 84(3), 645–657.

Fursch, H., 1977. Coccinellidenausbeuten aus Libanon und dem Iran im Museum Genf mit Beschreibung neuer Scymnini-Arten (Col. Cocc.). *Revue Suisse de Zoologie*, 84(3), 645-657.

Fursch, H., 1977. Coccinellideausbeuten aus Libanon und dem iran im Museum Genf mit Beshreibung near Scymnini-Arten (Col. Cocc.) *Revue Suisse Zoology* 84(3), 645-657.

Fursch, H., 1981. Eine neue *Nephus*-Art aus Iran (Coleoptera, Coccinellidae). *Revue suisse de Zoologie* 88 (1), 137-139.

Fursch, H., 1981. Eine neue *Nephus*-Art aus Iran (Coleoptera, Coccinellidae). *Revue Suisse de Zoologie* 88 (1),137-139.

Fursch, H., 1987. Neue Coccinelliden aus Aethiopien. *Folia Entomologica Hungarica* 39-44.

Fursch, H., 1995. A new *Serangium*-species from Iran (Col.: Coccinellidae). *Nachrichtenblatt der bayerischen Entomologen* 44, 20-22.

Fursch, H., 1997. Coccinellidae (Coleoptera) aus Rwanda. Bonn. *Zoologische Beitrage* 47 (1-2), 13-29.

Ghanbari Kohyani, A., Razmjou, M., Bagheri, M., 2013. Study of ladybirds (Col.: Coccinellidae) in Lordegan district. *Journal of International Academic Research for Multidisciplinary* 1(11), 249-254.

Ghanbari, A., Sadeghi, S.E., Ladan Moghadam, A.R., Fakhredini, M., 2012. An investigation on predaceous coccinellid's fauna, their distribution and determining dominant species on shading trees and shrubs in green spaces and parks of 16th Tehran municipal zone. Proceeding of the 20<sup>th</sup> Iranian Plant Protection Congress, 25-28 Aug 2012, Shiraz, Iran, pp 179.

Gholami Moghaddam, S., Vahedi, H., Heydazade, A., Ebrahimi, S., 2014. Faunal Study of ladybirds (Col.: Coccinellidae) in Kermanshah province and five new record for Kermanshah Province. Proceeding of the 21<sup>th</sup> Iranian Plant Protection Congress, 23-26 Aug 2014, Orumieh, Iran, pp 421.

Gordon, R.D., 1972. The tribe Noviini in the New World. *Journal of the Washington Academy of Sciences* 62 (1), 23-31.

Gordon, R.D., 1985. The Coccinellidae of America North of Mexico. *Journal of the New York Entomological Society* 93(1), 1-912.

Hahshenas, A.R., Malkeshi, S.H., Bagheri, M.R., 2004. The fauna of Coccinellids in cereal aphids and investigation on population fluctuation of dominant species in Isfahan province. Proceeding of the 16<sup>th</sup> Iranian Plant Protection Congress, 28 Aug-1 Sep 2004, Tabriz, Iran, pp 126.

Hajizadeh, J., Jalali, J., Peiravi, H., 2002. Introduction part of fauna lady beetles (Col.: Coccinellidae) of Gilan province. *Journal of Agriculture Science and Natural Resource* 9 (4), 99-112

Haji-Zadeh, J., Jalali, J., Peyrovi, H., 1998. A part of Coccinellid (Col.: Coccinellidae) fauna of Guilan pronince. Proceeding of the 13<sup>th</sup> Iranian Plant Protection Congress, 3-8 Sep 1998, Karaj, Iran, pp 212.

Hamedi, N., Moharrampour, S., 2012. Overwintering strategy in predatory ladybird *Brumus octosignatus* (Coleoptera: Coccinellidae). Proceeding of the 20<sup>th</sup> Iranian Plant Protection Congress, 25-28 Aug 2012, Shiraz, Iran, pp 610.

Hamidi, E., 2013. Faunisitic study of ladybird beetles (Col., Coccinellidae) in Boujnord district, Northern Khorasan province. M.Sc Thesis. College of plant production. Gorgan University, 114 p.

Heidari, M., Copland, M. J.W., 1989. Biological control of the mealybug *Pseudococcus affinis* (Mask.) (=*obscurus* Essig) by the coccinellid predator *Nephus (Scymnus) reunionii* (Fursch). Proceeding of the 9<sup>th</sup> Iranian Plant Protection Congress, 8-13 Sep 1989, Mashhad, Iran, pp 28.

Hoebke, E.R., Wheeler, A.G., 1996. Adventive lady beetles (Coleoptera: Coccinellidae) in the Canadian Maritime provinces. *Entomological News* 107, 281-290.

Iablokoff-Khanzorian, S.M, 1971. Synopsis des *Hyperaspis* paléarctiques (Col. Coccinellidae). Annales de la Société Entomologique de France 7(1), 163-200.

Iablokoff-Khanzorian, S.M., 1982. Les Coccinelles. Coléoptères-Coccinellidae. In: Société Nouvelles ed. Boubée press, Paris, 568 pp.

- Iablokoff-Khnzorian, S.M., 1974. Monographie der Gattung *Lithophilus* Frohlich (Col. Coccinellidae). Entomologische Arb Mus Georg Frey 25, 148-243.
- Iperti, G., 1999. Biodiversity of predaceous Coccinellidae in relation to bioindication and economic importance. Agriculture, Ecosystems & Environment 74, 323–342.
- Jafari, M., Sadeghi, E., Babalmorad, M., 2002. Natural enemies of poplar lace bug, *Monosteria unicostata* Mulstant & Rey (Tingidae) in Isfahan Province. Proceeding of the 15<sup>th</sup> Iranian Plant Protection Congress, 7-11 Sep 2002, Kermanshah, Iran, pp 120.
- Jafari, R., Fursch, H., Zare Khormizi, M., 2013. A checklist of the Scymninae (Coleoptera: Coccinellidae) of Iran. *International Research Journal of Applied and Basic Sciences* 5 (2), 154-160.
- Jafari, R., Kamali, K., 2007. Faunistic study of lady beetles (Col.: Coccinellidae) in Lorestan province and new records from Iran. *New Findings in Agriculture* 1 (4), 349-359.
- Jafari, R., Kamali, K., Shojai, M., Ostovan, H., 2008. The faunistic survey of ladybirds (Coleoptera, Coccinellidae) in Lorestan Province, Iran. Proceeding of the 18<sup>th</sup> Iranian Plant Protection Congress, 24-27 Aug 2008, Hamedan, Iran, pp 101.
- Jafari, R., Zarei, J.N., Vafaei, S.R., 2008. The faunistic survey on Coccinellids in Zarand Zone. *Journal of Entomological Research* 3(4), 277-284.
- Jalilvand, K., 2011. Preliminary survey on parasitoids of scale insect of Kermanshah city. Faculty of agriculture. M.Sc Thesis, Razi University, 180 pp.
- Jalilvand, K., Shirazi, M., Fallahzadeh, M., Ali Vahedi, H., Samih, M.A., Moeini Naghadeh, N., 2014. Survey of natural enemies of mealybug species (Hemiptera, Pseudococcidae) in Kermanshah province, Western Iran to inform biological control research. *Journal of the Entomological Research Society* 16(3), 01-10.
- Kalanrati, A., Sadeghi A., 2000. Investigation survey of ladybirds and determination of prevalent species in dry orchard almond in West Khorasan province. Proceeding of the 14<sup>th</sup> Iranian Plant Protection Congress, 4-7 Sep 2000, Isfahan, Iran, pp 271.
- Kaydan, M.B., Kilincer N., Uygun, N., Japoshvilli, G., S. Gaimari. 2006. Parasitoids and predators of Pseudococcidae (Hemiptera: Coccoidea) in Ankara, Turkey. *Phytoparasitica* 34(4), 331-337.
- Khalil Aria, A., Malkeshi, S.H., 2002. Identification of ladybirds of apple orchards in West Azerbaijan Province. Proceeding of the 15<sup>th</sup> Iranian Plant Protection Congress, 7-11 Sep 2002, Kermanshah, Iran, pp 88.
- Khandehroo, F., 2015. Survey on species diversity of arthropods associated with ash tree (*Fraxinus excelsior* L.) in Mashhad and vicinity. M.Sc Thesis. Ferdowsi University of Mashhad, 168p.
- Khormali, S., 2012. Biology of Twelve-spotted melon beetle, *Henosepilachna elaterii* in *Cucurbites* fields of Gonbad region. Proceeding of the 20<sup>th</sup> Iranian Plant Protection Congress, 25-28 Aug 2012, Shiraz, Iran, pp 676.

Khosravi, E., Vahedi, H., Mirmoayedi, A., 2014. Evaluation and identification predators of species *Trabutina mannipara* (Hemiptera: Pseudococcidae) in Kermanshah. Proceeding of the 21<sup>th</sup> Iranian Plant Protection Congress, 23-26 Aug 2014, Orumieh, Iran, pp 599.

Koohpayezadeh, K., Mossadegh, M.S., 1991. Some of the lady-birds (Coccinellidae) fauna of Kerman province. Proceeding of the 10<sup>th</sup> Iranian Plant Protection Congress, 1-5 Sep, Kerman, Iran, pp 64.

Korschefsky, R., 1931. PARS 118. Coccinellidae I. In: Coleopterorum catalogus. Schenklink S. (ed.): 225-659. Berlin; W. Junk.

Kovar I., 1995. Revision of the genera *Brumus* Muls. And *Exochomus* Redtb. (Coleoptera, Coccinellidae) of the Palaearctic Region. Part I. *Acta Entomologica Musei Nationalis Pragae*, 44.

Kovar, I., 2007. Coccinellidae, pp. 568-631. - In I. Löbl & A. Smetana (ed.): *Catalogue of Palaearctic Coleoptera*, Vol. 4. Stenstrup: Apollo Books, 935 pp.

Lachinani, P., Ahmadi, A.A., 1993. The natural enemies of yellow scale *Aonidiella orientalis* in citrus orchards of Fars province. Proceeding of the 11<sup>th</sup> Iranian Plant Protection Congress, 27 Aug-1 Sep 1993, Rasht, Iran, pp 196.

Lal, R., Kanakavalli, S., 1960. The genitalia of some Indian Coccinellidae. *The Annals of Zoology* 3(6), 69-110.

Larson, D.J., 2013. Key to Lady Beetles (Coleoptera: Coccinellidae) of Saskatchewan.

Löbl, I., Smetana, A., 2007. Catalogue of Palaearctic Coleoptera, Vol. 4. Stenstrup: Apollo Books, 935 pp.

Lotfalizadeh, H., 2001. Sexual discrimination in a part of fauna ladybirds of Moghan region. *Journal of entomological society of Iran* 21 (1), 69-88.

Lotfalizadeh, H., 2002. Natural enemies of cotton aphids in Moghan region, Northwest of Iran. Proceeding of the 15<sup>th</sup> Iranian Plant Protection Congress, 7-11 Sep 2002, Kermanshah, Iran, pp 36.

Lotfalizadeh, H., Hatami, B., Khalghani, J., 2000. Biological study *Exochomus quadripustulatus* (L.) (Col.: Coleoptera) on cypress tree mealybug, *Planococcus rosae* (Nasanov) (Hom.: Pseudococcidae) in Shiraz. Proceeding of the 14<sup>th</sup> Iranian Plant Protection Congress, 4-7 Sep 2000, Isfahan, Iran, pp 138.

Maafi, S.H., Rajabi, G.h., Jafari, M.E., 1998. Introduction of the natural enemies of *Planococcus citri* in Mazandaraan. Proceedings of the 13<sup>th</sup> Iranian Plant Protection Congress, 3-8 Sep 1998, Karaj, Iran, pp 153.

Mader, L., 1955. Evidenz der palaearktischen Coccinelliden und ihrer Aberrationen in Wort und Bild. II Teil. Entomologische Arbeiten aus dem Museum G. Frey Tutzing bei Muenchen 6, 764–1035.

Malkeshi, H., Rezwani, A., Talabi, A.A., 1998. Identification of important natural enemies of pome fruit trees aphids in Bojnord. Proceedings of the 13<sup>th</sup> Iranian Plant Protection Congress, 3-8 Sep 1998, Karaj, Iran, pp 163.

Merkviladze M., Kvavadze et al., 1985. Fauna i ekologija nekotorikh grupp nasekomikh i kleshchie grussi – Tbilisi, Metsniereba 3-27 (in Russian)

Modarres, S., Mossadegh, M.S., 1995. The coccinellids (Coleoptera) of southeast Khorasan province, Iran. Proceeding of the 12<sup>th</sup> Iranian Plant Protection Congress, 2-7 Sep 1995, Karaj, Iran, pp 326.

Modarres Awal, M., 1994. List of agricultural pests and their natural enemies in Iran. University of Ferdousi Publishers, Mashhad, 364 pp.

Modarres Najaf Abadi, S.S., Arjmand Nejad, A.R., Hanifian, S., 2008. The study on population fluctuation of cabbage aphid (*Brevicoryne brassicae*) and identification of its natural enemies in Sistan region. Proceeding of the 18<sup>th</sup> Iranian Plant Protection Congress, 24-27 Aug 2008, Hamedan, Iran, pp 38.

Modarres Najaf Abadi, S.S., Arjmand Nejad, A.R., Hejazi, R., 2008. Seasonal population changes of wheat green aphid (*Schizaphis graminum*) and introduction of its natural enemies in Sistan region. Proceeding of the 18<sup>th</sup> Iranian Plant Protection Congress, 24-27 Aug 2008, Hamedan, Iran, pp 353.

Mohammadbeigi, A., 2000. Natural enemies of the walnut aphids in Ghazvin region. Proceeding of the 14<sup>th</sup> Iranian Plant Protection Congress, 4-7 Sep 2000, Isfahan, Iran, pp 273.

Mohammadi, M., Hajizadeh, J., Lotfalizadeh, H., 2013. First report of *Vibidia duodecimguttata* (Poda, 1761) (Coleoptera: Coccinellidae) from Iran. Plant Pests Research 2(4), 68-70.

Mohhamad Poor, A., Jafari, R., Biravand, A., Zare, M., Rafiei Karahrudi, Z., 2013. Ladybirds associated with pomegranate trees in Lorestan province of Iran (Coleoptera: Coccinellidae). International Research Journal of Applied and Basic Sciences 5 (12), 1585-1589.

Mohhamad Poor, A., Jafari, R., Rafiei Karahrudi, Z., 2013. The faunistic survey of predatory ladybeetles (Coleoptera, Coccinellidae) in the Aleshtar region (Lorestan province), Iran. International Journal of Agriculture and Crop Sciences 6 (11), 723-728

Mohiseni, A.A., Keyhanian, A.A., Taghaddosi, M.V., Boroomand, H., 1998. The Coccinellids (Coleoptera) of olive orchads in Tarom-Olya, Zandjan. Proceeding of the 13<sup>th</sup> Iranian Plant Protection Congress, 3-8 Sep 1998, Karaj, Iran, pp 155.

Montazeri, M.M., Mossadegh, M.S., 1995. The coccinellids (Coleoptera) fauna of Gorgan plain and Gonbad Kavus. Proceedings of the 12<sup>th</sup> Iranian Plant Protection Congress, 2-7 Sep 1995, Karaj, Iran, pp 325.

Moodi, S., Mossadegh, M.S., 2000. The overwintering sites of *Brumus octosignatus* Gelber (Col.: Coccinellidae) in southeast Khorasan. Proceedings of the 14<sup>th</sup> Iranian Plant Protection Congress, 4-7 Sep 2000, Isfahan, Iran, pp 180.

Mosadegh, M.S., Aleosfoor, M., 2004. *Aphis nerii* B. de F. and its predatory natural enemies in Shiraz and Khuzestan. Proceeding of the 16<sup>th</sup> Iranian Plant Protection Congress, 28 Aug-1 Sep 2004, Tabriz, Iran, pp 43.

Mosadegh, M.S., Aleosfoor, M., 2004. Studies on the host preferences of *Aphis nerii* B. de F. lady birds. 16<sup>th</sup> Iranian Plant Protection Congress, 28 Aug-1 Sep 2004, Tabriz, Iran, pp 18.

Mossadegh, M.S., Eslamizadeh, R., Esfandiari, M., 2008. Biology study of mealybug *Nipaecoccus viridis* (New.) and possibility of its biological control by *Cryptolaemus montrouzieri* Mul. in citrus orchards of North Khuzestan. Proceeding of the 18<sup>th</sup> Iranian Plant Protection Congress, 24-27 Aug 2008, Hamedan, Iran, pp 35.

Mossadegh, M.S., Kocheili, F., 1993. Some of the lady-birds (Coccinellidae) fauna of Khuzestan, Iran. *Abvaz Science Journal of Agriculture* 16 (1,2), 23-30.

Mossadegh, M.S., Vafaei, S.H., Zarghami, S., Kocheli, F., Farsi, A., Alizadeh, M.S., Rezaei, N., 2012. Natural enemies of *Phenacoccus solenopsis* Tinsley (Sternorrhyncha: Coccoidea: Pseudococcidae) in Khuzestan, Iran. Proceedings of the 20<sup>th</sup> Iranian Plant Protection Congress, 25-28 Aug 2012, Shiraz, Iran, pp 216.

Naim, A., 1971. The fauna of Iranian Coccinellidae (1). *Entomologie et Phytopathologie Appliquees* 31, 11-14.

Nazari, A., Haji-Zadeh, J., Sahragard, A., 2000. Biology of *Exochomus nigromaculatus* Goez (Col.: Coccinellidae) in Guilan province. Proceedings of the 14<sup>th</sup> Iranian Plant Protection Congress, 4-7 Sep 2000, Isfahan, Iran, pp 322.

Nazari, A., Sahragard, A., Hajizadeh, J., 2006. Study on food consumption of *Exochomus nigromaculatus* Goeze (Coleoptera: Coccinellidae), and effect of temperature on larval food consumption. *Journal of Agriculture Sciences* 12 (1), 36-41.

Nedvěd O., Kovář I., 2012: Phylogeny and classification. In: Hodek I, van Emden H, Honěk A: Ecology and Behaviour of the Ladybird Beetles (Coccinellidae). Wiley-Blackwell, pp. 1–12.

Nematollahi, M.R., Bagheri, M.R., 2002. Part of coccinellid fauna on damask rose in Kashan region. Proceedings of the 15<sup>th</sup> Iranian Plant Protection Congress, 7-11 Sep 2002, Kermanshah, Iran, pp 176.

Novin, M., Mossadegh, M.S., Karami Nwjad, M., Ghasemi Nejad, M., 2000. Natural enemy of *Nipaecoccus viridis* (Newstead) in the north of Khuzestan. Proceedings of the 14<sup>th</sup> Iranian Plant Protection Congress, 4-7 Sep 2000, Isfahan, Iran, pp 264.

Omid, R., Esmaili, M., Sadeghi, E., 2000. *Chilocorus bipustulatus* L. (Col. Coccinellidae), the most important natural enemy of pine scale, *Leucaspis pusilla* Loew. In Tehran province. Proceeding of the 14<sup>th</sup> Iranian Plant Protection Congress, 4-7 Sep 2000, Isfahan, Iran, pp 295.

- Omid, R., Raoufi, B., Shrifinia, M., 1993. Investigation on biology and natural enemies of *Galerucella luteola* (Muller) (Col.: Chrysomelidae) in Tehran. Proceeding of the 11<sup>th</sup> Iranian Plant Protection Congress, 27 Aug-1 Sep 1993, Rasht, Iran, pp 243.
- Parvizi, R., Mostaan, M., Boromand, H., Mirzayans, H., 1986. A higher survey on fauna ladybirds of Western Azerbaijan. *Journal of Plant pests and deseases* 54 (1,2), 159-162.
- Plaza Infante, E., 1986. Contribución al conocimiento de los Coccinellidae españoles. Subfamilia Chilocorinae y Sticholotinae (Col.). *Eos* 62, 247-269.
- Poorani, J., 2004. An annotated checklist of the Coccinellidae (Coleoptera) of Indian subregion .Available from: [http://www.angelfire.com/bug2/j\\_poorani/](http://www.angelfire.com/bug2/j_poorani/) (accessed 5 March 2014).
- Portakaldali, M., Satar, S., 2010. Research of Coccinellidae (Coleoptera) fauna in Artvine and Rize province. Bitki Koruma Bulteni 50(3), 89-99.
- Pour, A.A., Shakarami, J., 2012. Recognition of ladybird fauna (Col.: Coccinellidae) in the alfalfa fields of Khoramabad. *Journal of Animal and Plant Sciences* 22(4), 939-943.
- Raimundo, A., 1992. Novas species de Scymnini para de fauna de coccinelideos de Portugal. Acta do V congress Iberico de Entomologia. Suplemento no 3. *Bol. Soc. Port. Entomol* 3, 373-384
- Raimundo, A.A.C., Fursch, H., van Harten, A., 2006. Notes on the ladybird beetles (Coleoptera: Coccinellidae) of Yemen, with the description of two new species. Fauna of Arabia, Frankfurt a. Main 21, 217-245.
- Raimundo, A.C., Fursch, H., van Harten, A., 2008, Order Coleoptera, family Coccinellidae. In: van Harten, A., (Ed.). Arthropod Fauna of the United Arab Emirates. Dar Al Ummah printing. Adu Dhabi, United Arab Emirates 1, 217-239.
- Raimundo, A.C., van Harten, A., 2000. An annotated checklist of the Coccinellidae (Insecta: Coleoptera) of Yemen. *Fauna of Arabia* 18, 211-243
- Razmjou, J., Hajizadeh, J., 2000. The coccinellids fauna of cotton fields in Moghan Region. Proceeding of the 14<sup>th</sup> Iranian Plant Protection Congress, 4-7 Sep 2000, Isfahan, Iran, pp 240.
- Rebolledo, R., Palma, R., Klein, C., Aguilera, Y.A., 2007. Coccinellini (Col. Coccinellidae) presentes en diferentes estratos vegetales en la IX Región de La Araucanía (Chile). *Idesia* 25, 63-71.
- Sadaghian, B., Nikdel, M., Dordaei, A.A., 2000. Faunistic study on Coleoptera order in Arasbaran. Proceedings of the 14<sup>th</sup> Iranian Plant Protection Congress, 4-7 Sep 2000, Isfahan, Iran, pp 303.
- Sadeghi, S.E., Ahmadi, S.M., Ebrahimi, E., 2002. Natural enemisies of alder leaf beetle *Galerucella lineola* F. (Col. Chrysomelidae) in Golestan Province. Proceeding of the 15<sup>th</sup> Iranian Plant Protection Congress, 7-11 Sep 2002, Kermanshah, Iran, pp 116.
- Sadeghi, S.E., Khanjani, M., 1998. A study of Coccinellidae fauna in alfa alfa fields in Hamedan. Proceedings of the 13<sup>th</sup> Iranian Plant Protection Congress, 3-8 Sep 1998, Karaj, Iran, pp 56.

- Saeid, H., 1995. Biology of *Epilachna chrysomelina* F. (Col., Coccinellidae) in laboratory. Proceedings of the 12<sup>th</sup> Iranian Plant Protection Congress, 2-7 Sep 1995, Karaj, Iran, pp 151.
- Saeedi, K., 1998. The Coccinellidae fauna of alfalfa fields in Boyer Ahmad regio. Proceedings of the 13<sup>th</sup> Iranian Plant Protection Congress, 3-8 Sep 1998, Karaj, Iran, pp 55.
- Saeedi, K., 2002. Preliminary studies of natural enemies of the Alfalfa Weevil, in Yasouj region. Proceedings of the 15<sup>th</sup> Iranian Plant Protection Congress, 7-11 Sep 2002, Kermanshah, Iran, pp 27.
- Salehi, T., Mehrnejad, M.R., Pashaei Rad, S., 2013. Diversity pattern of adult ladybird (Coleoptera: Coccinellidae) communities on pistachio trees in southern parts of Iran in different months. *Zoology and Ecology* 23(4), 286-292.
- Salehi, T., Pashaei Rad, S.H., Mehrnejad, M.R., Shokri, M.R., 2011. Ladybirds associated with pistachio trees in part of Kerman Province, Iran (Coleoptera: Coccinellidae). *Iranian Journal of Animal Biosystematics* 7(2), 157-169.
- Samin, N., Shojai, M., 2013. A study on Coccinellidae (Coleoptera: Cucujoidea) from Varamin and vicinity, Iran. *Linzer Biol. Beitr* 45(2), 2121-2126.
- Sasaji, H., 1971. Fauna Japonica: Coccinellidae (Ins. Col.). Tokyo, 340 pp.
- Korschefsky R., 1932. Coccinellidae II. In: W. Junk & S. Schenkling (Eds), Coleopterorum Catalogus. Part 120. Junk, Berlin, 435 pp.
- Seifollahi, A.R., Ebadi, R., Sadeghi, S.E., 2000. Natural enemies of Gaz psyllid (*Cyamophila dicora* Lovinova) in Isfahan Province. Proceeding of the 14<sup>th</sup> Iranian Plant Protection Congress, 4-7 Sep 2000, Isfahan, Iran, pp 299.
- Vandenberg, N.J., 2002. Coccinellidae Latreille 1807. In: Arnett, R. H.Jr., Thomas, M. C., Skelley, P. E., & Frank, J. H. ed. American Beetles 2, Polyphaga: Scarabaeoidea through Curculionoidea. CRC Press, Boca Raton, New York, pp. 371–389.
- Vodjdani, S., 1965. Useful and pest Coccinellids of Iran. University of Tehran publishers, Karaj, 101 pp.
- Yaghmaee, F., Kharazi, A., 1995. A faunistic survey of coccinellids in Mashhad region. Proceedings of the 12<sup>th</sup> Iranian Plant Protection Congress, 2-7 Sep 1995, Karaj, Iran, pp 307.
- Yazdani, A., 1990. Fauna of Coccinellidae in Fars province. M.S.c Thessis, University of Kerman 71, 1-140.
- Yazdani, A., Ahmadi, A.A., 1991. New record of eight Coccinellid species of genus *Scymnus* from Fars province. Proceeding of the 10<sup>th</sup> Iranian Plant Protection Congress, 1-5 Sep, Kerman, Iran, pp 85.
- Yazdani, A., Ahmadi, A.A., 1992. Four coccinellid species of subfamily Sticholotidinae in Fars (Coleoptera: Coccinellidae). *Journal of Entomological Society of Iran* 11(1/2), 13-18.

Yazdani, A., Ebrahimik, J., 1993. Predators of scale insects on pistachio trees in the Kerman Province. Proceedings of the 11<sup>th</sup> Iranian Plant Protection Congress, 27 Aug-1 Sep 1993, Rasht, Iran, pp 210.

Yousof, D.E., Mahmoud, M.E.E., Mohamed, A.H., 2013. Prospects of biological control of daten palm green pit scale insect *Asterolecanium phoenicis* Rao (Homoptera: Asterolecaniidae) in Sudan. *Persian Gulf crop protection* 2(2), 42-48.

Zare Khormizi, M., Biranvand, A., Shakarami, J., 2013. The faunistic survey of lady beetles (Coleoptera: Coccinellidae) in the Mehriz region (Yazd Province), Iran. *Bulletin of Iraq Natural History Museum* 12(4), 43-51.

Zare Khormizi, M., Ostovan, H., Fallahzadeh, M., 2014. Status of ladybirds (Coleoptera: Coccinellidae) in Khatam County (Yazd Province) and the first report of *Hyperaspis reppensis* (Herbst, 1783) for Iranian fauna. *Journal of Biological Control* 28(3), 133-136.

Zare Khormizi, M., Ostovan, H., Fallahzadeh, M., Mossadegh, M.S., 2014. Report of three ladybird beetles (Coleoptera: Coccinellidae) from Iran. Proceeding of the 21<sup>th</sup> Iranian Plant Protection Congress, 23-26 Aug 2014, Orumieh, Iran, pp 490.

Zare, M., Jafari, R., Dehghan Dehnavi, L., Javan, S., 2012. Polymorphism of *Adalia bipunctata* Mulsant (Col.: Coccinellidae) in Center of Iran. *Global Advanced Research Journal of Microbiology* 1(2), 027-032.